

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
14 November 2002 (14.11.2002)

PCT

(10) International Publication Number
WO 02/090580 A1

(51) International Patent Classification⁷: **C12Q 1/68**

[MY/SG]; 4 Tan Boon Chong Avenue, 1027 Singapore (SG).

(21) International Application Number: PCT/AU02/00558

(22) International Filing Date: 3 May 2002 (03.05.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/288,441 3 May 2001 (03.05.2001) US

(71) Applicants (*for all designated States except US*): **NATIONAL CANCER CENTRE OF SINGAPORE PTE LTD** [SG/SG]; 11 Hospital Drive, 169610 Singapore (SG). **SINGAPORE GENERAL HOSPITAL PTE LTD** [SG/SG]; 11 Third Hospital, SNEC Building, 7th Floor, 168751 Singapore (SG).

(71) Applicant (*for AU only*): **ARGAET, Victor, Peter** [AU/AU]; 29 Beaconsfield Terrace, Gordon Park, Queensland 4031 (AU).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **HUYNH, The Hung** [CA/SG]; 1 Pearlbank #26-03, 169016 Singapore (SG). **CHOW, Pierce, Kah, Hoe** [MY/SG]; 10A Siak Kew Avenue, 348081 Singapore (SG). **SOO, Khee, Chee**

(74) Agents: **ARGAET, Victor, Peter** et al.; Davies Collison Cave, Level 3, 303 Coronation Drive, Milton, Queensland 4064 (AU).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: IDENTIFYING LIVER CANCER BY DETECTING ABERRANT EXPRESSION OF INSULIN LIKE GROWTH FACTOR BINDING PROTEIN

(57) Abstract: The invention is the linkage of the aberrant expression of insulin like growth factor binding proteins (IGFBPs) to the development and/or progression of liver cancer. It includes a method for detecting the presence or diagnosing the risk of a liver cancer, either before or after the onset of clinical symptoms, by detecting an aberrant level and/or functional activity of an expression product of a gene encoding an IGFBP, which correlates with the presence or risk of liver cancer. Also disclosed are methods of screening for agents that modulate the level and/or functional activity of an expression product of an IGFBP gene or gene belonging to the same biosynthetic or regulatory pathway as an IGFBP gene for use in methods for modulating liver cell proliferation of tumourgenesis, or for treating and/or preventing a liver cancer or related condition.



WO 02/090580 A1

**IDENTIFYING LIVER CANCER BY DETECTING
ABERRANT EXPRESSION OF INSULIN LIKE
GROWTH FACTOR BINDING PROTEIN**

FIELD OF THE INVENTION

THIS INVENTION relates generally to polynucleotides and polypeptides linked to cancer. More particularly, the present invention relates to the aberrant expression of insulin-like growth factor binding proteins (IGFBPs), which is linked to the development and/or progression of cancer, especially a cancer of the liver such as hepatocellular carcinoma (HCC). Even more particularly, the present invention relates to a method for detecting the presence or diagnosing the risk of a liver cancer, either before or after the onset of clinical symptoms, by detecting an aberrant level and/or functional activity of an expression product of a gene encoding an insulin-like growth factor binding protein (IGFBP), which correlates with the presence or risk of liver cancer. Assessing the level and/or functional activity of the aberrant expression product is useful as a prognostic indicator of disease outcome. The invention also encompasses methods of screening for agents that modulate the level and/or functional activity of an expression product of an *IGFBP* gene or a gene belonging to the same biosynthetic or regulatory pathway as an *IGFBP* gene for use in methods for modulating liver cell proliferation or tumorigenesis, of for treating and/or preventing a liver cancer or related condition.

Bibliographic details of the publications numerically referred to in this specification are collected at the end of the description.

BACKGROUND OF THE INVENTION

Hepatocellular carcinoma (HCC) is one of the most common malignancies worldwide but treatment outcomes for HCC have remained generally poor. The majority of patients with HCC have inoperable disease with very poor prognosis (1). Survival in patients with curative resection carried out at dedicated centres is between 35 - 50% at 5 years and much lower elsewhere (2, 3). Long-term survival is uncommon because of the frequent presence of recurrence, metastasis or the development of new primaries (4, 5). There is also currently no accepted adjuvant or palliative treatment modalities that have been conclusively shown to prolong survival in HCC (6).

Recent research has shown that insulin-like growth factors I and II (IGF-I and IGF-II) are potent mitogens for human hepatoma cells (7), and that both IGF-I mRNA (8) and IGF-I receptors (IGF-IR) (7, 8) are detectable in human hepatoma cell lines. Both IGF-

- 2 -

I and IGF-II bind with high affinity to specific IGF-binding proteins (IGFBPs), which modulate their bioactivity. At least six IGFBPs have been described (reviewed in (9, 10)). Expression of genes encoding the various IGFBPs has been observed in many tissues, and is subject to intricate physiological regulation (reviewed in (9, 10)). IGFBP-3 is the most
5 abundant IGF binding protein in the circulation, where it forms a 150 kDa complex with an acid-labile subunit and IGF-I or IGF-II (10). The IGFBP-3 gene is expressed in many tissues and IGFBP-3 has affinities for IGFs that are either equal to or stronger than those of the IGF receptors and therefore inhibits the IGFs by sequestration in the extracellular compartment (reviewed in (9, 10)). Recent evidence also demonstrates that IGFBP-3 has
10 growth inhibitory activity that is independent of its IGF binding properties (reviewed in (10)).

Existing evidence suggests that early diagnosis of HCC may allow modulation of its activities and the potential for therapeutical applications in the control of HCC. Conventional methods for diagnosis of HCC are based on measuring serum levels of
15 alpha-fetoprotein (AFP). Normally, AFP is a major serum protein synthesised by foetal liver cells, yolk sac cells, and in trace amounts by the foetal gastrointestinal tract. Reappearance of AFP in adult serum often signals pathologic conditions, particularly the presence of HCC and germ cell tumours containing yolk sac cell elements. Although existing assays may be used successfully for monitoring treatment of AFP-producing
20 tumours and as an independent prognostic tool, the finding of elevated serum AFP levels in some patients with nonmalignant liver diseases, particularly in acute and chronic viral hepatitis and cirrhosis, has limited the value of such assays as an independent specific test to establish the diagnosis of cancer.

An acute need, therefore, still exists in the art for an assay which is selective for
25 patients with HCC, particularly given the need for early detection of HCC in high risk populations.

SUMMARY OF THE INVENTION

The present invention is predicated in part on the determination that IGFBPs are differentially expressed in liver cancer (*e.g.*, HCC). Thus, IGFBPs and the transcripts from which they are translated have utility, for example, as markers of liver cancer, and as
5 targets for therapeutic intervention in liver cancer. Accordingly, in one aspect of the present invention, there is provided a method for detecting the presence or diagnosing the risk of a liver cancer in a patient, comprising detecting in a biological sample obtained from said patient aberrant expression of a gene encoding an insulin-like growth factor binding protein (IGFBP). Preferably, the IGFBP is selected from IGFBP-1, IGFBP-2 or
10 IGFBP-3. The liver cancer is preferably a liver cell carcinoma.

In another aspect, the invention contemplates a method for detecting the presence or diagnosing the risk of a liver cancer in a patient, comprising detecting in a biological sample obtained from said patient aberrant expression of at least one gene encoding an IGFBP selected from IGFBP-1, -2 and -3.

15 In yet another aspect, the invention encompasses a method for detecting the presence or diagnosing the risk of a liver cancer in a patient, comprising detecting in a biological sample obtained from said patient a level and/or functional activity of an expression product of a gene encoding an IGFBP-1, which is elevated relative to a normal reference level and/or functional activity of said expression product.

20 In still yet another aspect, the invention envisions a method for detecting the presence or diagnosing the risk of a liver cancer in a patient, comprising detecting in a biological sample obtained from said patient a level and/or functional activity of an expression product of a gene encoding an IGFBP-2, which is elevated relative to a normal reference level and/or functional activity of said expression product.

25 In a further aspect, the invention contemplates a method for detecting the presence or diagnosing the risk of a liver cancer in a patient, comprising detecting in a biological sample obtained from said patient a level and/or functional activity of an expression product of a gene encoding an IGFBP-3, which is reduced relative to a normal reference level and/or functional activity of said expression product.

30 In yet a further aspect, the invention resides in a method for detecting the presence or diagnosing the risk of a liver cancer in a patient, comprising detecting in a biological sample obtained from said patient:

- 4 -

– a level and/or functional activity of an expression product of an *IGFBP-1* gene, which is elevated relative to a normal reference level and/or functional activity of said *IGFBP-1* expression product; and/or

– a level and/or functional activity of an expression product of an *IGFBP-2* gene, which is elevated relative to a normal reference level and/or functional activity of said *IGFBP-2* expression product; and/or

– a level and/or functional activity of an expression product of an *IGFBP-3* gene, which is reduced relative to a normal reference level and/or functional activity of said *IGFBP-3* expression product.

In another aspect, the invention contemplates a method for detecting the presence or diagnosing the risk of a liver cancer in a patient, comprising detecting in a biological sample obtained from said patient:

– a level and/or functional activity of an expression product of an *IGFBP-1* gene, which is elevated relative to a normal reference level and/or functional activity of said *IGFBP-1* expression product; and

– a level and/or functional activity of an expression product of an *IGFBP-2* gene, which is elevated relative to a normal reference level and/or functional activity of said *IGFBP-2* expression product.

In yet another aspect, the invention encompasses a method for detecting the presence or diagnosing the risk of a liver cancer in a patient, comprising detecting in a biological sample obtained from said patient:

– a level and/or functional activity of an expression product of an *IGFBP-1* gene, which is elevated relative to a normal reference level and/or functional activity of said *IGFBP-1* expression product; and

– a level and/or functional activity of an expression product of an *IGFBP-3* gene, which is reduced relative to a normal reference level and/or functional activity of said *IGFBP-3* expression product.

In yet another aspect, the invention envisions a method for detecting the presence or diagnosing the risk of a liver cancer in a patient, comprising detecting in a biological sample obtained from said patient:

– a level and/or functional activity of an expression product of an *IGFBP-2* gene, which is elevated relative to a normal reference level and/or functional activity of said *IGFBP-2* expression product; and

- 5 -

– a level and/or functional activity of an expression product of an *IGFBP-3* gene, which is reduced relative to a normal reference level and/or functional activity of said *IGFBP-3* expression product.

In still yet another aspect, the invention features a method for detecting the presence or diagnosing the risk of a liver cancer in a patient, comprising detecting in a biological sample obtained from said patient:

– a level and/or functional activity of an expression product of an *IGFBP-1* gene, which is elevated relative to a normal reference level and/or functional activity of said *IGFBP-1* expression product; and

– a level and/or functional activity of an expression product of an *IGFBP-2* gene, which is elevated relative to a normal reference level and/or functional activity of said *IGFBP-2* expression product; and

– a level and/or functional activity of an expression product of an *IGFBP-3* gene, which is reduced relative to a normal reference level and/or functional activity of said *IGFBP-3* expression product.

In another aspect, the invention provides a method for diagnosis in a patient of the progression of a liver cancer, comprising detecting aberrant expression of at least one gene encoding an IGFBP in a biological sample obtained from said patient.

In yet another aspect, the invention contemplates a method for prognostic assessment of a liver cancer in a patient, comprising detecting in a biological sample obtained from said patient aberrant expression of at least one gene encoding an IGFBP.

In yet another aspect, the invention extends to the use of an agent in the manufacture of a medicament for treating and/or preventing a liver cancer, wherein said agent is optionally formulated with a pharmaceutically acceptable carrier and is identifiable by a screening assay comprising:

– contacting a preparation comprising at least a portion of an expression product of a gene selected from an *IGFBP* gene or a gene relating to the same regulatory or biosynthetic pathway as the *IGFBP* gene, or a variant or derivative of said expression product, or comprising a genetic sequence which regulates expression of said expression product and which is operably connected to a marker gene, with said agent; and

- 6 -

– detecting a change in the level and/or functional activity of said at least a portion of said expression product or said variant or derivative or a change in the level of expression of said marker gene.

In another aspect, the invention contemplates the use of an agent that reduces the level and/or functional activity of an expression product of a gene encoding an IGFBP—1 in the manufacture of a medicament for treating and/or preventing a liver cancer, wherein said agent is optionally formulated with a pharmaceutically acceptable carrier and is identifiable by a screening assay comprising:

– contacting a preparation comprising at least a portion of an expression product of an *IGFBP—1* gene, or a variant or derivative of said expression product, or comprising a genetic sequence which regulates expression of said expression product and which is operably connected to a marker gene, with said agent; and

– detecting a reduction in the level and/or functional activity of said at least a portion of said expression product or said variant or derivative or a reduction in the level and/or functional activity of an expression product of said marker gene.

In yet another aspect, the invention provides the use of an agent that reduces the level and/or functional activity of an expression product of a gene encoding an IGFBP—2 in the manufacture of a medicament for treating and/or preventing a liver cancer, wherein said agent is optionally formulated with a pharmaceutically acceptable carrier and is identifiable by a screening assay comprising:

– contacting a preparation comprising at least a portion of an expression product of an *IGFBP—2* gene, or a variant or derivative of said expression product, or comprising a genetic sequence which regulates expression of said expression product and which is operably connected to a marker gene, with said agent; and

– detecting a reduction in the level and/or functional activity of said at least a portion of said expression product or said variant or derivative or a reduction in the level and/or functional activity of an expression product of said marker gene.

In a further aspect, the invention extends to the use of an agent that increases the level and/or functional activity of an expression product of a gene encoding an IGFBP—3 in the manufacture of a medicament for treating and/or preventing a liver cancer, wherein said agent is optionally formulated with a pharmaceutically acceptable carrier and is identifiable by a screening assay comprising:

- 7 -

– contacting a preparation comprising at least a portion of an expression product of an *IGFBP-3* gene, or a variant or derivative of said expression product, or comprising a genetic sequence which regulates expression of said expression product and which is operably connected to a marker gene, with said agent; and

5 – detecting an increase in the level and/or functional activity of said at least a portion of said expression product or said variant or derivative or an increase in the level and/or functional activity of an expression product of said marker gene.

In one embodiment, the agent is selected from (1) an antisense oligonucleotide or ribozyme that binds to, or otherwise interacts specifically with, a polynucleotide selected
10 from an *IGFBP-1* polynucleotide, an *IGFBP-2* polynucleotide or complement of these, or (2) an antigen-binding molecule that is immuno-interactive with IGFBP-1 and/or IGFBP-2. In another embodiment, the agent is selected from a first member comprising an IGFBP-3 polypeptide or a biologically active fragment thereof, or variant or derivative of these or a second member comprising a polynucleotide encoding said first member, which
15 is operably connected to a regulatory polynucleotide.

In yet another aspect, the invention contemplates the use of a vector in the manufacture of a medicament, which is optionally formulated with a pharmaceutically acceptable carrier, for the treatment and/or prophylaxis of a liver cancer or related condition, wherein said vector comprises at least a portion of an *IGFBP-3* polynucleotide
20 operably linked, in the sense direction, to a regulatory polynucleotide.

In another aspect, the invention extends to the use of a vector in the manufacture of a medicament, which is optionally formulated with a pharmaceutically acceptable carrier, for the treatment and/or prophylaxis of a liver cancer or related condition, wherein said vector comprises an isolated *IGFBP-1* polynucleotide operably linked, in the
25 antisense direction, to a regulatory polynucleotide.

In yet another aspect, the invention provides the use of a vector in the manufacture of a medicament, which is optionally formulated with a pharmaceutically acceptable carrier, for the treatment and/or prophylaxis of a liver cancer or related condition, wherein said vector comprises an isolated *IGFBP-2* polynucleotide operably
30 linked, in the antisense direction, to a regulatory polynucleotide.

In still another aspect, the invention contemplates a method for modulating liver cell proliferation or tumorigenesis, comprising contacting a liver cell with an agent as

- 8 -

broadly described above for a time and under conditions sufficient to modulate the level and/or functional activity of an *IGFBP* expression product.

According to another aspect, the invention provides a method for treatment and/or prophylaxis of a liver cancer or related condition, comprising administering to a patient in
5 need of such treatment an effective amount of a compound selected from an agent as broadly described above, an agent-containing medicament as broadly described above, or a vector-containing medicament as broadly described above.

The invention also contemplates, in another aspect, the use of an agent-containing medicament as broadly described above, or a vector-containing composition as broadly
10 described above in the preparation of a medicament for treating a liver cancer or related condition.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1. IGFBP-3 expression in adjacent normal liver tissue and HCC tumours. Tissue lysates from normal adjacent liver tissue and HCC tumours were analysed by Western blotting. Blots were incubated with anti-human IGFBP-3 and α -tubulin antibodies. Representative samples are shown in panels (A), (B) and (C). All normal adjacent liver tissues had high levels of IGFBP-3 while IGFBP-3 protein was either under-expressed or absent from HCC tumours. Note that the low molecular weight IGFBP-3 fragments were present in serum samples (see panel (C)).

Figure 2. Immunostaining of normal liver (A) and HCC tumours (B) for IGFBP-3. Normal adjacent liver tissue and HCC tumours were stained with anti-human IGFBP-3 as described in Example 1. Adjacent normal liver tissue, showing intense expression of normal cells for IGFBP-3 while very low staining signal was observed for the morphologically disorganised HCC cells. Representative staining are shown. (original magnification X 400).

Figure 3. Immunostaining of normal liver and HCC tumours for IGF-I and IGF-II. Normal adjacent liver tissue (AC) and HCC tumours (B,D) were stained with anti-human IGF-I (A and B) or anti-human IGF-II (C and D) as described in Example 1. Representative samples are shown. Normal adjacent liver tissue shows intense staining for IGF-I while low staining signals are seen for HCC cells. HCC tumours show more intense staining for IGF-II than normal adjacent tissue. (original magnification X 400).

Figure 4. IGF-I receptor expression in adjacent normal liver tissue and HCC tumours. Tissue lysates from normal adjacent liver tissue and HCC tumours were analysed by Western blotting as described in Example 1. Blots were incubated with anti-IGF-IR (A) and anti- α -tubulin (B) antibodies. Note that in certain pairs of samples, normal adjacent liver tissue had higher levels of IGF-IR than HCC tumours.

Figure 5. Effects of IGFs, IGFBP-3, and IGF-II antibody on HepG2 cell proliferation. Cells were cultured as described in Example 1. Cells were incubated with serum free MEM, and indicated reagents of IGF-I for 48 h. Cell number was determined as described (14). Means of triplicate experiments were plotted. Bars with different letters are significantly different from one another at ($p < 0.01$, Mann Whitney U test). SE at each point was $< 15\%$. *A*, Effect of IGF-I on HepG2 proliferation. IGF-I significantly increased HepG2 cell number. *B*, Effect of IGFBP-3 on HepG2 cell proliferation. Growth of HepG2

- 10 -

cells was significantly inhibited by exogenous IGFBP-3. *C*, Effect of hrIGFBP-3 on IGF-I-induced HepG2 cell proliferation. Cells were incubated with either serum free medium or 25 ng/mL IGF-I in the presence of various concentrations of hrIGFBP-3 for 48 h. IGF-I-induced HepG2 proliferation was significantly attenuated by hrIGFBP-3. *D*, Effect of IGF-II and anti-IGF-II antibody on HepG2 cell proliferation. Cells were incubated with serum free, 25 ng/mL hrIGF-II, rabbit preimmune serum (1:800 dilution), rabbit anti-human IGF-II antibody (2 μ g/mL) and 25 ng/mL IGF-I in combination with rabbit anti-human IGF-II (2 μ g/mL). IGF-II-induced proliferation rates were significantly decreased in the presence of anti-IGF-II antibody. Note that IGF-II antibody also inhibits HepG2 cell growth.

Figure 6. Serum levels of IGFBP-1, -2, -3 and alpha fetoprotein in non-HCC and HCC patients. Serum samples from non-HCC and HCC patients were analysed by Western blotting as described in Example 2. Blots were incubated with anti-human IGFBP-1, anti-bovine IGFBP-2, anti-human IGFBP-3 and anti-human alpha fetoprotein antibodies. Representative samples are shown in panels 6A – 6P. All non-HCC serum samples had high levels of IGFBP-3 while IGFBP-1 and IGFBP-2 were low. IGFBP-3 was either low or absent from HCC. Upregulation of IGFBP-1 and IGFBP-2 was observed in most HCC serum samples. Note that AFP was absent in some HCC serum samples.

BRIEF DESCRIPTION OF THE SEQUENCES: SUMMARY TABLE**TABLE A**

<i>SEQUENCE ID</i>	<i>DESCRIPTION</i>	<i>LENGTH</i>
SEQ ID NO: 1	Nucleotide sequence corresponding to human insulin-like growth factor binding protein 1 (IGFBP-1), mRNA as set forth in GenBank Accession No. XM_004688	1514 nts
SEQ ID NO: 2	IGFBP-1 polypeptide encoded by SEQ ID NO: 1	259 aa
SEQ ID NO: 3	Nucleotide sequence corresponding to human insulin-like growth factor binding protein 2 (IGFBP-2), mRNA as set forth in GenBank Accession No. NM_000597	1433 nts
SEQ ID NO: 4	IGFBP-2 polypeptide encoded by SEQ ID NO: 3	328 aa
SEQ ID NO: 5	Nucleotide sequence corresponding to human insulin-like growth factor binding protein 3 (IGFBP-3), mRNA as set forth in GenBank Accession No. BC000013	2497 nts
SEQ ID NO: 6.	IGFBP-3 polypeptide encoded by SEQ ID NO: 5	291 aa
SEQ ID NO: 7	Nucleotide sequence corresponding to human insulin-like growth factor binding protein 1 (IGFBP-1) gene as set forth in GenBank Accession No. M74587	6128 nts
SEQ ID NO: 8	IGFBP-1 polypeptide encoded by SEQ ID NO: 7	259 aa
SEQ ID NO: 9	Nucleotide sequence corresponding to human insulin-like growth factor binding protein 2 (IGFBP-2) gene as set forth in GenBank Accession No. gi 18552832:211494-242517	31024 nts
SEQ ID NO: 10	IGFBP-2 polypeptide encoded by SEQ ID NO: 7	328 aa

- 12 -

<i>SEQUENCE ID</i>	<i>DESCRIPTION</i>	<i>LENGTH</i>
SEQ ID NO: 11	Nucleotide sequence corresponding to human insulin-like growth factor binding protein 3 (IGFBP-3) gene as set forth in GenBank Accession No. M35878	10884 nts
SEQ ID NO: 12	IGFBP-3 polypeptide encoded by SEQ ID NO: 11	291 aa

DETAILED DESCRIPTION OF THE INVENTION

1. Definitions

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by those of ordinary skill in the art to which the invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, preferred methods and materials are described. For the purposes of the present invention, the following terms are defined below.

The articles “a” and “an” are used herein to refer to one or to more than one (i.e. to at least one) of the grammatical object of the article. By way of example, “an element” means one element or more than one element.

The term “*aberrant expression product*” refers to expressed polynucleotides or polypeptide, which result from a substitution, deletion and/or addition of one or more nucleotides or amino acid residues in a “normal” reference polynucleotide or polypeptide, and which correlate with the presence or risk of a liver cancer.

“*Amplification product*” refers to a nucleic acid product generated by nucleic acid amplification techniques.

By “*antigen-binding molecule*” is meant a molecule that has binding affinity for a target antigen. It will be understood that this term extends to immunoglobulins, immunoglobulin fragments and non-immunoglobulin derived protein frameworks that exhibit antigen-binding activity.

The term “*biological sample*” as used herein refers to a sample that may be extracted, untreated, treated, diluted or concentrated from a patient. The biological sample may comprise a fluid selected from whole blood, serum, plasma, saliva, urine, sweat, ascitic fluid, peritoneal fluid, synovial fluid, amniotic fluid, cerebrospinal fluid, tissue biopsy (e.g., liver biopsy), and the like. The biological sample preferably includes serum, whole blood, plasma and lymph or other circulatory fluid, saliva, mucus secretion and respiratory fluid. More preferably, the biological sample comprises a circulatory fluid such as serum or whole blood or a fractionated portion thereof. Most preferably, the biological sample comprises serum or a fractionated portion thereof.

- 14 -

By “*biologically active fragment*” is meant a fragment of a full-length parent polypeptide which fragment retains the activity of the parent polypeptide. A biologically active fragment will therefore have, for example, the activity of an IGFBP. As used herein, the term “*biologically active fragment*” includes deletion variants and small peptides, for example of at least 10, preferably at least 20 and more preferably at least 30 contiguous amino acids, which comprise the above activities. Peptides of this type may be obtained through the application of standard recombinant nucleic acid techniques or synthesised using conventional liquid or solid phase synthesis techniques. For example, reference may be made to solution synthesis or solid phase synthesis as described, for example, in Chapter 9 entitled “*Peptide Synthesis*” by Atherton and Shephard which is included in a publication entitled “*Synthetic Vaccines*” edited by Nicholson and published by Blackwell Scientific Publications. Alternatively, peptides can be produced by digestion of a polypeptide of the invention with proteinases such as endoLys-C, endoArg-C, endoGlu-C and staphylococcus V8-protease. The digested fragments can be purified by, for example, high performance liquid chromatographic (HPLC) techniques.

Throughout this specification, unless the context requires otherwise, the words “*comprise*”, “*comprises*” and “*comprising*” will be understood to imply the inclusion of a stated step or element or group of steps or elements but not the exclusion of any other step or element or group of steps or elements.

By “*effective amount*”, in the context of treating or preventing a condition is meant the administration of that amount of active to an individual in need of such treatment or prophylaxis, either in a single dose or as part of a series, that is effective for treatment or prophylaxis of that condition. The effective amount will vary depending upon the health and physical condition of the individual to be treated, the taxonomic group of individual to be treated, the formulation of the composition, the assessment of the medical situation, and other relevant factors. It is expected that the amount will fall in a relatively broad range that can be determined through routine trials.

As used herein, the term “*function*” refers to a biological, enzymatic, or therapeutic function.

Reference herein to “*immuno-interactive*” includes reference to any interaction, reaction, or other form of association between molecules and in particular where one of the molecules is, or mimics, a component of the immune system.

- 15 -

By “*isolated*” is meant material that is substantially or essentially free from components that normally accompany it in its native state.

By “*marker gene*” is meant a gene that imparts a distinct phenotype to cells expressing the marker gene and thus allows such transformed cells to be distinguished from cells that do not have the marker. A selectable marker gene confers a trait for which one can ‘select’ based on resistance to a selective agent (*e.g.*, antibiotic, radiation, heat, or other treatment damaging to untransformed cells). A screenable marker gene (or marker gene) confers a trait that one can identify through observation or testing, *i.e.*, by ‘screening’ (*e.g.* β -glucuronidase, luciferase, or other enzyme activity not present in untransformed cells).

By “*obtained from*” is meant that a sample such as, for example, a polynucleotide extract or polypeptide extract is isolated from, or derived from, a particular source of the host. For example, the extract can be obtained from a tissue or a biological fluid isolated directly from the host.

The term “*oligonucleotide*” as used herein refers to a polymer composed of a multiplicity of nucleotide residues (deoxyribonucleotides or ribonucleotides, or related structural variants or synthetic analogues thereof) linked via phosphodiester bonds (or related structural variants or synthetic analogues thereof). Thus, while the term “oligonucleotide” typically refers to a nucleotide polymer in which the nucleotide residues and linkages between them are naturally occurring, it will be understood that the term also includes within its scope various analogues including, but not restricted to, peptide nucleic acids (PNAs), phosphoramidates, phosphorothioates, methyl phosphonates, 2-O-methyl ribonucleic acids, and the like. The exact size of the molecule can vary depending on the particular application. An oligonucleotide is typically rather short in length, generally from about 10 to 30 nucleotide residues, but the term can refer to molecules of any length, although the term “polynucleotide” or “nucleic acid” is typically used for large oligonucleotides.

The term “*operably connected*” or “*operably linked*” as used herein means placing a structural gene under the regulatory control of a promoter, which then controls the transcription and optionally translation of the gene. In the construction of heterologous promoter/structural gene combinations, it is generally preferred to position the genetic sequence or promoter at a distance from the gene transcription start site that is approximately the same as the distance between that genetic sequence or promoter and the

- 16 -

gene it controls in its natural setting; *i.e.* the gene from which the genetic sequence or promoter is derived. As is known in the art, some variation in this distance can be accommodated without loss of function. Similarly, the preferred positioning of a regulatory sequence element with respect to a heterologous gene to be placed under its control is defined by the positioning of the element in its natural setting; *i.e.* the genes from which it is derived.

The term “*patient*” refers to patients of human or other mammal and includes any individual it is desired to examine or treat using the methods of the invention. However, it will be understood that “*patient*” does not imply that symptoms are present. Suitable mammals that fall within the scope of the invention include, but are not restricted to, primates, livestock animals (*eg.* sheep, cows, horses, donkeys, pigs), laboratory test animals (*eg.* rabbits, mice, rats, guinea pigs, hamsters), companion animals (*eg.* cats, dogs) and captive wild animals (*eg.* foxes, deer, dingoes).

By “*pharmaceutically acceptable carrier*” is meant a solid or liquid filler, diluent or encapsulating substance that can be safely used in topical or systemic administration to a animal, preferably a mammal including humans.

The term “*polynucleotide*” or “*nucleic acid*” as used herein designates mRNA, RNA, cRNA, cDNA or DNA. The term typically refers to oligonucleotides greater than 30 nucleotide residues in length. Polynucleotide sequences are understood to encompass complementary strands as well as alternative backbones described herein.

The terms “*polynucleotide variant*” and “*variant*” refer to polynucleotides displaying substantial sequence identity with a reference polynucleotide sequence or polynucleotides that hybridise with a reference sequence under stringent conditions that are defined hereinafter. These terms also encompasses polynucleotides in which one or more nucleotides have been added or deleted, or replaced with different nucleotides. In this regard, it is well understood in the art that certain alterations inclusive of mutations, additions, deletions and substitutions can be made to a reference polynucleotide whereby the altered polynucleotide retains the biological function or activity of the reference polynucleotide. The terms “*polynucleotide variant*” and “*variant*” also include naturally occurring allelic variants.

“*Polypeptide*”, “*peptide*” and “*protein*” are used interchangeably herein to refer to a polymer of amino acid residues and to variants and synthetic analogues of the same. Thus, these terms apply to amino acid polymers in which one or more amino acid residues

- 17 -

is a synthetic non-naturally occurring amino acid, such as a chemical analogue of a corresponding naturally occurring amino acid, as well as to naturally-occurring amino acid polymers.

The term “*polypeptide variant*” refers to polypeptides in which one or more amino acids have been replaced by different amino acids. It is well understood in the art that some amino acids may be changed to others with broadly similar properties without changing the nature of the activity of the polypeptide (conservative substitutions).

By “*primer*” is meant an oligonucleotide which, when paired with a strand of DNA, is capable of initiating the synthesis of a primer extension product in the presence of a suitable polymerising agent. The primer is preferably single-stranded for maximum efficiency in amplification but can alternatively be double-stranded. A primer must be sufficiently long to prime the synthesis of extension products in the presence of the polymerisation agent. The length of the primer depends on many factors, including application, temperature to be employed, template reaction conditions, other reagents, and source of primers. For example, depending on the complexity of the target sequence, the oligonucleotide primer typically contains 15 to 35 or more nucleotide residues, although it can contain fewer nucleotide residues. Primers can be large polynucleotides, such as from about 200 nucleotide residues to several kilobases or more. Primers can be selected to be “substantially complementary” to the sequence on the template to which it is designed to hybridise and serve as a site for the initiation of synthesis. By “substantially complementary”, it is meant that the primer is sufficiently complementary to hybridise with a target polynucleotide. Preferably, the primer contains no mismatches with the template to which it is designed to hybridise but this is not essential. For example, non-complementary nucleotide residues can be attached to the 5' end of the primer, with the remainder of the primer sequence being complementary to the template. Alternatively, non-complementary nucleotide residues or a stretch of non-complementary nucleotide residues can be interspersed into a primer, provided that the primer sequence has sufficient complementarity with the sequence of the template to hybridise therewith and thereby form a template for synthesis of the extension product of the primer.

“*Probe*” refers to a molecule that binds to a specific sequence or sub-sequence or other moiety of another molecule. Unless otherwise indicated, the term “probe” typically refers to a polynucleotide probe that binds to another polynucleotide, often called the “target polynucleotide”, through complementary base pairing. Probes can bind target

polynucleotides lacking complete sequence complementarity with the probe, depending on the stringency of the hybridisation conditions. Probes can be labelled directly or indirectly.

The term “*recombinant polynucleotide*” as used herein refers to a polynucleotide formed *in vitro* by the manipulation of a polynucleotide into a form not normally found in nature. For example, the recombinant polynucleotide can be in the form of an expression vector. Generally, such expression vectors include transcriptional and translational regulatory polynucleotide operably linked to the polynucleotide.

By “*recombinant polypeptide*” is meant a polypeptide made using recombinant techniques, *ie.* through the expression of a recombinant or synthetic polynucleotide.

By “*reporter molecule*” as used in the present specification is meant a molecule that, by its chemical nature, provides an analytically identifiable signal that allows the detection of a complex comprising an antigen-binding molecule and its target antigen. The term “reporter molecule” also extends to use of cell agglutination or inhibition of agglutination such as red blood cells on latex beads, and the like.

By “*vector*” is meant a polynucleotide molecule, preferably a DNA molecule derived, for example, from a plasmid, bacteriophage, yeast or virus, into which a polynucleotide can be inserted or cloned. A vector preferably contains one or more unique restriction sites and can be capable of autonomous replication in a defined host cell including a target cell or tissue or a progenitor cell or tissue thereof, or be integrable with the genome of the defined host such that the cloned sequence is reproducible. Accordingly, the vector can be an autonomously replicating vector, *ie.* a vector that exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, *eg.* a linear or closed circular plasmid, an extrachromosomal element, a minichromosome, or an artificial chromosome. The vector can contain any means for assuring self-replication. Alternatively, the vector can be one which, when introduced into the host cell, is integrated into the genome and replicated together with the chromosome(s) into which it has been integrated. A vector system can comprise a single vector or plasmid, two or more vectors or plasmids, which together contain the total DNA to be introduced into the genome of the host cell, or a transposon. The choice of the vector will typically depend on the compatibility of the vector with the host cell into which the vector is to be introduced. In the present case, the vector is preferably a viral or viral-derived vector, which is operably functional in animal and preferably mammalian cells. Such vector may be derived from a poxvirus, an adenovirus or yeast. The vector can also include a selection

- 19 -

marker such as an antibiotic resistance gene that can be used for selection of suitable transformants. Examples of such resistance genes are known to those of skill in the art and include the *nptII* gene that confers resistance to the antibiotics kanamycin and G418 (Geneticin®) and the *hph* gene which confers resistance to the antibiotic hygromycin B.

5 As used herein, underscoring or italicising the name of a gene shall indicate the gene, in contrast to its protein product, which is indicated in the absence of any underscoring or italicising. For example, “*IGFBP-1*” shall mean the *IGFBP-1* gene, whereas “IGFBP-1” shall indicate the protein product of the “*IGFBP-1*” gene.

2. *Methods of detecting aberrant IGFBP expression*

10 The present invention is predicated in part on the discovery that the sera of patients with liver cancer, particularly hepatocellular carcinoma, have aberrant levels of insulin-like growth factor binding proteins (IGFBPs) relative to sera of normal patients. Thus, the invention features a method for detecting the presence or diagnosing the risk of a liver cancer in a patient, comprising detecting aberrant expression of a gene encoding an
15 IGFBP in a biological sample obtained from said patient. The aberrant expression preferably relates to a level and/or functional activity of an expression product of said gene, which correlates with the presence or risk of said liver cancer. A correlation with the presence or risk of a liver cancer is made for example when the level and/or functional activity of an *IGFBP* expression product in the biological sample differs by at least 10%,
20 more preferably at least 50%, even more preferably at least 100%, even more preferably at least 200%, even more preferably at least 400%, even more preferably at least 600% and still even more preferably at least 1000% compared to the level and/or functional activity of said expression product in a biological sample obtained from normal patients or from patients who are not afflicted with that liver cancer. In a preferred embodiment, the level
25 and/or functional activity of an *IGFBP-1* or *IGFBP-2* expression product in said biological sample is at least 10%, more preferably at least 50%, even more preferably at least 100%, even more preferably at least 200%, even more preferably at least 400%, even more preferably at least 600% and still even more preferably at least 1000% higher than the level and/or functional activity of that expression product in a biological sample
30 obtained from normal patients or from patients who are not afflicted with said liver cancer. In another preferred embodiment, the level and/or functional activity of an *IGFBP-3* expression product in said biological sample is at least 10%, more preferably at least 50%, even more preferably at least 100%, even more preferably at least 200%, even more

- 20 -

preferably at least 400%, even more preferably at least 600% and still even more preferably at least 1000% lower than the level and/or functional activity of that expression product in a biological sample obtained from normal patients or from patients who are not afflicted with said liver cancer.

5 Accordingly, the presence or risk of liver cancer is diagnosed when (1) an expression product of *IGFBP-1* or *IGFBP-2* is expressed at a higher level, or (2) an expression product of *IGFBP-3* is expressed at a lower level, compared to the level at which it is expressed in normal patients or in patients without liver cancer. In a preferred embodiment of this type, the method comprises detecting a level and/or functional activity
10 of an expression product of an *IGFBP-1* gene or an *IGFBP-2* gene, which is elevated relative to a normal reference level and/or functional activity of that expression product. In another preferred embodiment of this type, the method comprises detecting a level and/or functional activity of an expression product of an *IGFBP-3* gene, which is reduced relative to a normal reference level and/or functional activity of that expression product.

15 Thus, it may be desirable for example to qualitatively or quantitatively determine IGFBP protein levels and/or *IGFBP* transcription levels. Alternatively or additionally, it may be desirable to search for aberrant *IGFBP* structural genes and regulatory regions, which encode aberrant *IGFBP* expression products or which produce aberrant levels of *IGFBP* expression products. Alternatively or additionally, it may be desirable to
20 qualitatively or quantitatively determine the level of an expression product (e.g., transcript, protein) of a gene relating to the same regulatory or biosynthetic pathway as an *IGFBP* gene, which can modulate or otherwise influence IGFBP protein levels and/or *IGFBP* transcription levels. Likewise, it may also be desirable to search for an aberrant gene relating to the same regulatory or biosynthetic pathway as an *IGFBP* gene.

25 The biological sample can be any tissue (e.g., a liver biopsy) or fluid. The biological sample is preferably a biological fluid including, but not restricted to, a circulatory fluid such as whole blood, serum, plasma and lymph as well as other circulatory fluid and saliva, mucus secretion and respiratory fluid. More preferably, the biological sample is serum or a fractionated portion thereof.

30 2.1 Genetic Diagnosis

One embodiment of the instant invention comprises a method for detecting an increase in the expression of an *IGFBP* gene by qualitatively or quantitatively determining

- 21 -

the transcripts of an *IGFBP* gene in a cell (*e.g.*, a liver cell such a Kupffer cell). Exemplary human nucleic acid sequences for *IGFBP-1*, *IGFBP-2* and *IGFBP-3* mRNA and there corresponding genes are set forth in the enclosed Sequence Listing *infra* and are summarised in TABLE A *supra*.

5 Another embodiment of the instant invention comprises a method for detecting reduction in the expression or function of an *IGFBP* gene (*e.g.*, *IGFBP-3*), or enhancement of expression or function of an *IGFBP* gene (*e.g.*, *IGFBP-1*, *IGFBP-2*), by examining the genes and transcripts of a cell. Nucleic acid used is isolated from cells contained in the biological sample, according to standard methodologies (Sambrook, *et al.*, "Molecular
10 Cloning. A Laboratory Manual", Cold Spring Harbor Press, 1989; Ausubel *et al.*, "Current Protocols in Molecular Biology", John Wiley & Sons Inc, 1994-1998). The nucleic acid may be genomic DNA or fractionated or whole cell RNA. Where RNA is used, it may be desired to convert the RNA to a complementary DNA. In one embodiment, the RNA is whole cell RNA; in another, it is poly-A RNA. In one embodiment, the nucleic acid is
15 amplified by a nucleic acid amplification technique. Suitable nucleic acid amplification techniques are well known to the skilled artisan, and include the polymerase chain reaction (PCR) as for example described in Ausubel *et al.* (*supra*); strand displacement amplification (SDA) as for example described in U.S. Patent No 5,422,252; rolling circle replication (RCR) as for example described in Liu *et al.*, (1996) and International
20 application WO 92/01813) and Lizardi *et al.*, (International Application WO 97/19193); nucleic acid sequence-based amplification (NASBA) as for example described by Sooknanan *et al.*, (1994, *Biotechniques* 17:1077-1080); and Q- β replicase amplification as for example described by Tyagi *et al.*, (1996, *Proc. Natl. Acad. Sci. USA* 93: 5395-5400).

 Depending on the format, the specific nucleic acid of interest is identified in the
25 sample directly using amplification or with a second, known nucleic acid following amplification. Next, the identified product is detected. In certain applications, the detection may be performed by visual means (*e.g.*, ethidium bromide staining of a gel). Alternatively, the detection may involve indirect identification of the product via chemiluminescence, radioactive scintigraphy of radiolabel or fluorescent label or even via
30 a system using electrical or thermal impulse signals (Affymax Technology; Bellus, 1994, *J Macromol. Sci. Pure, Appl. Chem.*, A31(1): 1355-1376).

 Following detection, one may compare the results seen in a given patient with a control reaction or a statistically significant reference group of normal patients. In this

- 22 -

way, it is possible to correlate the amount of an IGFBP detected with the progression or severity of the disease.

In addition to determining levels of *IGFBP* transcripts, it also may prove useful to examine various types of defects. These defect could include deletions, insertions, point mutations and duplications. Point mutations result in stop codons, frameshift mutations or amino acid substitutions. Somatic mutations are those occurring in non-germline tissues. Germ-line tissue can occur in any tissue and are inherited. Mutations in and outside the coding region also may affect the amount of IGFBP produced, both by altering the transcription of the gene or in destabilising or otherwise altering the processing of either the transcript (mRNA) or protein.

A variety of different assays are contemplated in this regard, including but not limited to, fluorescent *in situ* hybridisation (FISH), direct DNA sequencing, pulse field gel electrophoresis (PFGE) analysis, Southern or Northern blotting, single-stranded conformation analysis (SSCA), RNase protection assay, allele-specific oligonucleotide (ASO), dot blot analysis, denaturing gradient gel electrophoresis, RFLP and PCR-SSCP.

2.1.1 Primers and Probes

Primers may be provided in double-stranded or single-stranded form, although the single-stranded form is preferred. Probes, while perhaps capable of priming, are designed to bind to a target DNA or RNA and need not be used in an amplification process. In preferred embodiments, the probes or primers are labelled with radioactive species ^{32}P , ^{14}C , ^{35}S , ^3H , or other label), with a fluorophore (rhodamine, fluorescein) or a chemilluminiscent label (luciferase).

2.1.2 Template Dependent Amplification Methods

A number of template dependent processes are available to amplify the target sequences present in a given template sample. An exemplary nucleic acid amplification technique is the polymerase chain reaction (referred to as PCR) which is described in detail in U.S. Pat. Nos. 4,683,195, 4,683,202 and 4,800,159, Ausubel *et al.* (*supra*), and in Innis *et al.*, ("PCR Protocols", Academic Press, Inc., San Diego Calif., 1990).

Briefly, in PCR, two primer sequences are prepared that are complementary to regions on opposite complementary strands of the target sequence. An excess of deoxynucleoside triphosphates are added to a reaction mixture along with a DNA polymerase, *e.g.*, Taq polymerase. If the target sequence is present in a sample, the primers

- 23 -

will bind to the target and the polymerase will cause the primers to be extended along the target sequence by adding on nucleotides. By raising and lowering the temperature of the reaction mixture, the extended primers will dissociate from the target to form reaction products, excess primers will bind to the target and to the reaction products and the process is repeated.

A reverse transcriptase PCR amplification procedure may be performed in order to quantify the amount of mRNA amplified. Methods of reverse transcribing RNA into cDNA are well known and described in Sambrook *et al.*, 1989. Alternative methods for reverse transcription utilise thermostable, RNA-dependent DNA polymerases. These methods are described in WO 90/07641. Polymerase chain reaction methodologies are well known in the art.

Another method for amplification is the ligase chain reaction ("LCR"), disclosed in EPO No. 320 308. In LCR, two complementary probe pairs are prepared, and in the presence of the target sequence, each pair will bind to opposite complementary strands of the target such that they abut. In the presence of a ligase, the two probe pairs will link to form a single unit. By temperature cycling, as in PCR, bound ligated units dissociate from the target and then serve as "target sequences" for ligation of excess probe pairs. U.S. Pat. No. 4,883,750 describes a method similar to LCR for binding probe pairs to a target sequence.

Q β Replicase, described in PCT Application No. PCT/US87/00880, may also be used as still another amplification method in the present invention. In this method, a replicative sequence of RNA that has a region complementary to that of a target is added to a sample in the presence of an RNA polymerase. The polymerase will copy the replicative sequence that can then be detected.

An isothermal amplification method, in which restriction endonucleases and ligases are used to achieve the amplification of target molecules that contain nucleotide 5' α -thio-triphosphates in one strand of a restriction site may also be useful in the amplification of nucleic acids in the present invention, Walker *et al.*, (1992, Proc. Natl. Acad. Sci. U.S.A 89: 392-396).

Strand Displacement Amplification (SDA) is another method of carrying out isothermal amplification of nucleic acids which involves multiple rounds of strand displacement and synthesis, i.e., nick translation. A similar method, called Repair Chain Reaction (RCR), involves annealing several probes throughout a region targeted for

- 24 -

amplification, followed by a repair reaction in which only two of the four bases are present. The other two bases can be added as biotinylated derivatives for easy detection. A similar approach is used in SDA. Target specific sequences can also be detected using a cyclic probe reaction (CPR). In CPR, a probe having 3' and 5' sequences of non-specific DNA and a middle sequence of specific RNA is hybridised to DNA that is present in a sample. Upon hybridisation, the reaction is treated with RNase H, and the products of the probe identified as distinctive products that are released after digestion. The original template is annealed to another cycling probe and the reaction is repeated.

Still another amplification method, described in GB Application No. 2 202 328, and in PCT Application No. PCT/US89/01025, may be used in accordance with the present invention. In the former application, "modified" primers are used in a PCR-like, template- and enzyme-dependent synthesis. The primers may be modified by labelling with a capture moiety (*e.g.*, biotin) and/or a detector moiety (*e.g.*, enzyme). In the latter application, an excess of labelled probes are added to a sample. In the presence of the target sequence, the probe binds and is cleaved catalytically. After cleavage, the target sequence is released intact to be bound by excess probe. Cleavage of the labelled probe signals the presence of the target sequence.

Other nucleic acid amplification procedures include transcription-based amplification systems (TAS), including nucleic acid sequence based amplification (NASBA) and 3SR (Kwoh *et al.*, 1989, *Proc. Natl. Acad. Sci. U.S.A.*, **86**: 1173; Gingeras *et al.*, PCT Application WO 88/10315). In NASBA, the nucleic acids can be prepared for amplification by standard phenol/chloroform extraction, heat denaturation of a clinical sample, treatment with lysis buffer and minispin columns for isolation of DNA and RNA or guanidinium chloride extraction of RNA. These amplification techniques involve annealing a primer which has target specific sequences. Following polymerisation, DNA/RNA hybrids are digested with RNase H while double stranded DNA molecules are heat denatured again. In either case the single stranded DNA is made fully double stranded by addition of second target specific primer, followed by polymerisation. The double-stranded DNA molecules are then multiply transcribed by an RNA polymerase such as T7 or SP6. In an isothermal cyclic reaction, the RNAs are reverse transcribed into single stranded DNA, which is then converted to double stranded DNA, and then transcribed once again with an RNA polymerase such as T7 or SP6. The resulting products, whether truncated or complete, indicate target specific sequences.

- 25 -

Davey *et al.*, EPO No. 329 822 disclose a nucleic acid amplification process involving cyclically synthesising single-stranded RNA ("ssRNA"), ssDNA, and double-stranded DNA (dsDNA), which may be used in accordance with the present invention. The ssRNA is a template for a first primer oligonucleotide, which is elongated by reverse transcriptase (RNA-dependent DNA polymerase). The RNA is then removed from the resulting DNA:RNA duplex by the action of ribonuclease H (RNase H, an RNase specific for RNA in duplex with either DNA or RNA). The resultant ssDNA is a template for a second primer, which also includes the sequences of an RNA polymerase promoter (exemplified by T7 RNA polymerase) 5' to its homology to the template. This primer is then extended by DNA polymerase (exemplified by the large "Klenow" fragment of *E. coli* DNA polymerase I), resulting in a double-stranded DNA ("dsDNA") molecule, having a sequence identical to that of the original RNA between the primers and having additionally, at one end, a promoter sequence. This promoter sequence can be used by the appropriate RNA polymerase to make many RNA copies of the DNA. These copies can then re-enter the cycle leading to very swift amplification. With proper choice of enzymes, this amplification can be done isothermally without addition of enzymes at each cycle. Because of the cyclical nature of this process, the starting sequence can be chosen to be in the form of either DNA or RNA.

Miller *et al.* in PCT Application WO 89/06700 disclose a nucleic acid sequence amplification scheme based on the hybridisation of a promoter/primer sequence to a target single-stranded DNA ("ssDNA") followed by transcription of many RNA copies of the sequence. This scheme is not cyclic, *i.e.*, new templates are not produced from the resultant RNA transcripts. Other amplification methods include "RACE" and "one-sided PCR" (Frohman, M. A., In: "PCR Protocols: A Guide to Methods and Applications", Academic Press, N.Y., 1990; Ohara *et al.*, 1989, *Proc. Natl Acad. Sci. U.S.A.*, **86**: 5673-5677).

Methods based on ligation of two (or more) oligonucleotides in the presence of nucleic acid having the sequence of the resulting "di-oligonucleotide", thereby amplifying the di-oligonucleotide, may also be used in the amplification step of the present invention. Wu *et al.*, (1989, *Genomics* **4**: 560).

2.1.3 Southern/Northern Blotting

Blotting techniques are well known to those of skill in the art. Southern blotting involves the use of DNA as a target, whereas Northern blotting involves the use of RNA as

- 26 -

a target. Each provide different types of information, although cDNA blotting is analogous, in many aspects, to blotting or RNA species.

Briefly, a probe is used to target a DNA or RNA species that has been immobilized on a suitable matrix, often a filter of nitrocellulose. The different species
5 should be spatially separated to facilitate analysis. This often is accomplished by gel electrophoresis of nucleic acid species followed by "blotting" on to the filter.

Subsequently, the blotted target is incubated with a probe (usually labelled) under conditions that promote denaturation and rehybridization. Because the probe is designed to base pair with the target, the probe will binding a portion of the target sequence under
10 renaturing conditions. Unbound probe is then removed, and detection is accomplished as described above.

2.1.4 Detection Methods

Products may be visualised in order to confirm amplification of the target sequences. One typical visualisation method involves staining of a gel with ethidium
15 bromide and visualisation under UV light. Alternatively, if the amplification products are integrally labelled with radio- or fluorometrically-labelled nucleotides, the amplification products can then be exposed to x-ray film or visualised under the appropriate stimulating spectra, following separation.

In one embodiment, visualisation is achieved indirectly. Following separation of
20 amplification products, a labelled nucleic acid probe is brought into contact with the amplified target sequence. The probe preferably is conjugated to a chromophore but may be radiolabelled. In another embodiment, the probe is conjugated to a binding partner, such as an antibody or biotin, and the other member of the binding pair carries a detectable moiety or reporter molecule.

25 In one embodiment, detection is by a labelled probe. The techniques involved are well known to those of skill in the art and can be found in many standard texts on molecular protocols. See Sambrook *et al.*, 1989. For example, chromophore or radiolabel probes or primers identify the target during or following amplification.

One example of the foregoing is described in U.S. Pat. No. 5,279,721, which
30 discloses an apparatus and method for the automated electrophoresis and transfer of nucleic acids. The apparatus permits electrophoresis and blotting without external

- 27 -

manipulation of the gel and is ideally suited to carrying out methods according to the present invention.

In addition, the amplification products described above may be subjected to sequence analysis to identify specific kinds of variations using standard sequence analysis techniques. Within certain methods, exhaustive analysis of genes is carried out by sequence analysis using primer sets designed for optimal sequencing (Pignon *et al.*, 1994, *Hum. Mutat.* 3: 126-132). The present invention provides methods by which any or all of these types of analyses may be used. Using, for example, the sequences set forth herein, oligonucleotide primers may be designed to permit the amplification of sequences throughout *IGFBP-1*, *-2*, and *-3* genes that may then be analysed by direct sequencing.

2.1.5 Kit Components

All the essential materials and reagents required for detecting and sequencing *IGFBP* genes and variants thereof may be assembled together in a kit. The kits may also optionally include appropriate reagents for detection of labels, positive and negative controls, washing solutions, dilution buffers and the like. For example, a nucleic acid-based detection kit may include (i) a polynucleotide according to the invention (which may be used as a positive control), (ii) an oligonucleotide primer according to the invention. Also included may be enzymes suitable for amplifying nucleic acids including various polymerases (Reverse Transcriptase, Taq, Sequenase™ DNA ligase etc. depending on the nucleic acid amplification technique employed), deoxynucleotides and buffers to provide the necessary reaction mixture for amplification. Such kits also generally will comprise, in suitable means, distinct containers for each individual reagent and enzyme as well as for each primer or probe. The kit may also contain instructions for performing the method of the invention.

2.1.6 Chip Technologies

Also contemplated by the present invention are chip-based DNA technologies such as those described by Hacia *et al.* (1996, *Nature Genetics* 14: 441-447) and Shoemaker *et al.* (1996, *Nature Genetics* 14: 450-456). Briefly, these techniques involve quantitative methods for analysing large numbers of genes rapidly and accurately. By tagging genes with oligonucleotides or using fixed probe arrays, one can employ chip technology to segregate target molecules as high density arrays and screen these molecules

- 28 -

on the basis of hybridisation. See also Pease *et al.* (1994, *Proc. Natl. Acad. Sci. U.S.A.* **91**: 5022-5026); Fodor *et al.* (1991, *Science* **251**: 767-773).

2.2 Protein-based diagnostics

2.2.1 Antigen-binding molecules

5 Antigen-binding molecules that are immuno-interactive with a target molecule of the present invention can be used in measuring an increase or decrease in *IGFBP* expression. Thus, the present invention also contemplates antigen-binding molecules that bind specifically to IGFBP-1, IGFBP-2 or IGFBP-3 polypeptides or to proteins that regulate or otherwise influence the level and/or functional activity of one or more of the
10 aforesaid IGFBP polypeptides. For example, the antigen-binding molecules may comprise whole polyclonal antibodies. Such antibodies may be prepared, for example, by injecting a target molecule of the invention into a production species, which may include mice or rabbits, to obtain polyclonal antisera. Methods of producing polyclonal antibodies are well known to those skilled in the art. Exemplary protocols which may be used are described for
15 example in Coligan *et al.*, "Current Protocols In Immunology", (John Wiley & Sons, Inc, 1991), and Ausubel *et al.*, (1994-1998, *supra*), in particular Section III of Chapter 11.

 In lieu of the polyclonal antisera obtained in the production species, monoclonal antibodies may be produced using the standard method as described, for example, by Köhler and Milstein (1975, *Nature* **256**, 495-497), or by more recent modifications thereof
20 as described, for example, in Coligan *et al.*, (1991, *supra*) by immortalising spleen or other antibody-producing cells derived from a production species which has been inoculated with target molecule of the invention. A number of human and other mammalian antibodies to IGFBP-1, -2 and -3 are known, and are available either commercially or through techniques well known to the art and industry.

25 The invention also contemplates as antigen-binding molecules Fv, Fab, Fab' and F(ab')₂ immunoglobulin fragments. Alternatively, the antigen-binding molecule may comprise a synthetic stabilised Fv fragment. Exemplary fragments of this type include single chain Fv fragments (sFv, frequently termed scFv) in which a peptide linker is used to bridge the N terminus or C terminus of a V_H domain with the C terminus or N-terminus,
30 respectively, of a V_L domain. ScFv lack all constant parts of whole antibodies and are not able to activate complement. Suitable peptide linkers for joining the V_H and V_L domains are those which allow the V_H and V_L domains to fold into a single polypeptide chain

- 29 -

having an antigen binding site with a three dimensional structure similar to that of the antigen binding site of a whole antibody from which the Fv fragment is derived. Linkers having the desired properties may be obtained by the method disclosed in U.S. Patent No 4,946,778. However, in some cases a linker is absent. ScFvs may be prepared, for example, in accordance with methods outlined in Kreber *et al* (Kreber *et al.* 1997, *J. Immunol. Methods*; **201**(1): 35-55). Alternatively, they may be prepared by methods described in U.S. Patent No 5,091,513, European Patent No 239,400 or the articles by Winter and Milstein (1991, *Nature* **349**:293) and Plünckthun *et al* (1996, In *Antibody engineering: A practical approach*. 203-252).

Alternatively, the synthetic stabilised Fv fragment comprises a disulphide stabilised Fv (dsFv) in which cysteine residues are introduced into the V_H and V_L domains such that in the fully folded Fv molecule the two residues will form a disulphide bond therebetween. Suitable methods of producing dsFv are described for example in (Glockscuther *et al.* *Biochem.* **29**: 1363-1367; Reiter *et al.* 1994, *J. Biol. Chem.* **269**: 18327-18331; Reiter *et al.* 1994, *Biochem.* **33**: 5451-5459; Reiter *et al.* 1994, *Cancer Res.* **54**: 2714-2718; Webber *et al.* 1995, *Mol. Immunol.* **32**: 249-258).

Also contemplated as antigen-binding molecules are single variable region domains (termed dAbs) as for example disclosed in (Ward *et al.* 1989, *Nature* **341**: 544-546; Hamers-Casterman *et al.* 1993, *Nature.* **363**: 446-448; Davies & Riechmann, 1994, *FEBS Lett.* **339**: 285-290).

Alternatively, the antigen-binding molecule may comprise a "minibody". In this regard, minibodies are small versions of whole antibodies, which encode in a single chain the essential elements of a whole antibody. Suitably, the minibody is comprised of the V_H and V_L domains of a native antibody fused to the hinge region and CH3 domain of the immunoglobulin molecule as, for example, disclosed in U.S. Patent No 5,837,821.

In an alternate embodiment, the antigen binding molecule may comprise non-immunoglobulin derived, protein frameworks. For example, reference may be made to (Ku & Schultz, 1995, *Proc. Natl. Acad. Sci. USA*, **92**: 652-6556) which discloses a four-helix bundle protein cytochrome b562 having two loops randomised to create complementarity determining regions (CDRs), which have been selected for antigen binding.

The antigen-binding molecule may be multivalent (*ie.* having more than one antigen-binding site). Such multivalent molecules may be specific for one or more antigens. Multivalent molecules of this type may be prepared by dimerisation of two

- 30 -

antibody fragments through a cysteinyl-containing peptide as, for example disclosed by (Adams *et al.*, 1993, *Cancer Res.* **53**: 4026-4034; Cumber *et al.*, 1992, *J. Immunol.* **149**: 120-126). Alternatively, dimerisation may be facilitated by fusion of the antibody fragments to amphiphilic helices that naturally dimerise (Pack P. Plünckthun, 1992, *Biochem.* **31**: 1579-1584), or by use of domains (such as the leucine zippers jun and fos) that preferentially heterodimerise (Kostelny *et al.*, 1992, *J. Immunol.* **148**: 1547-1553). In an alternate embodiment, the multivalent molecule may comprise a multivalent single chain antibody (multi-scFv) comprising at least two scFvs linked together by a peptide linker. In this regard, non-covalently or covalently linked scFv dimers termed “diabodies” may be used. Multi-scFvs may be bispecific or greater depending on the number of scFvs employed having different antigen binding specificities. Multi-scFvs may be prepared for example by methods disclosed in U.S. Patent No. 5,892,020.

The antigen-binding molecules of the invention may be used for affinity chromatography in isolating a natural or recombinant polypeptide or biologically active fragment of the invention. For example reference may be made to immunoaffinity chromatographic procedures described in Chapter 9.5 of Coligan *et al.*, (1995-1997, *supra*).

2.2.2 Immunodiagnostic assays

The above antigen-binding molecules have utility in measuring directly or indirectly modulation of *IGFBP* expression in healthy and diseased states, through techniques such as ELISAs and Western blotting. Illustrative assay strategies which can be used to detect a target polypeptide of the invention include, but are not limited to, immunoassays involving the binding of an antigen-binding molecule to the target polypeptide (*e.g.*, an *IGFBP* polypeptide) in the sample, and the detection of a complex comprising the antigen-binding molecule and the target polypeptide. Preferred immunoassays are those that can measure the level and/or functional activity of a target molecule of the invention. Typically, an antigen-binding molecule that is immuno-interactive with a target polypeptide of the invention is contacted with a biological sample suspected of containing said target polypeptide. The concentration of a complex comprising the antigen-binding molecule and the target polypeptide is measure in and the measured complex concentration is then related to the concentration of target polypeptide in the sample. Consistent with the present invention, the presence of an aberrant

- 31 -

concentration of the target polypeptide is indicative of the presence or risk of a liver cancer.

Any suitable technique for determining formation of an antigen-binding molecule-target antigen complex may be used. For example, an antigen-binding molecule according to the invention, having a reporter molecule associated therewith may be utilised in immunoassays. Such immunoassays include, but are not limited to, radioimmunoassays (RIAs), enzyme-linked immunosorbent assays (ELISAs) and immunochromatographic techniques (ICTs), Western blotting which are well known those of skill in the art. For example, reference may be made to Coligan *et al.* (1994, *supra*) which discloses a variety of immunoassays that may be used in accordance with the present invention. Immunoassays may include competitive assays as understood in the art or as for example described *infra*. It will be understood that the present invention encompasses qualitative and quantitative immunoassays.

Suitable immunoassay techniques are described for example in US Patent Nos. 4,016,043, 4, 424,279 and 4,018,653. These include both single-site and two-site assays of the non-competitive types, as well as the traditional competitive binding assays. These assays also include direct binding of a labelled antigen-binding molecule to a target antigen.

Two site assays are particularly favoured for use in the present invention. A number of variations of these assays exist, all of which are intended to be encompassed by the present invention. Briefly, in a typical forward assay, an unlabelled antigen-binding molecule such as an unlabelled antibody is immobilised on a solid substrate and the sample to be tested brought into contact with the bound molecule. After a suitable period of incubation, for a period of time sufficient to allow formation of an antibody-antigen complex, another antigen-binding molecule, suitably a second antibody specific to the antigen, labelled with a reporter molecule capable of producing a detectable signal is then added and incubated, allowing time sufficient for the formation of another complex of antibody-antigen-labelled antibody. Any unreacted material is washed away and the presence of the antigen is determined by observation of a signal produced by the reporter molecule. The results may be either qualitative, by simple observation of the visible signal, or may be quantitated by comparing with a control sample containing known amounts of antigen. Variations on the forward assay include a simultaneous assay, in which both sample and labelled antibody are added simultaneously to the bound antibody. These

- 32 -

techniques are well known to those skilled in the art, including minor variations as will be readily apparent. In accordance with the present invention, the sample is one that might contain an antigen including a tissue or fluid as described above.

In the typical forward assay, a first antibody having specificity for the antigen or antigenic parts thereof is either covalently or passively bound to a solid surface. The solid surface is typically glass or a polymer, the most commonly used polymers being cellulose, polyacrylamide, nylon, polystyrene, polyvinyl chloride or polypropylene. The solid supports may be in the form of tubes, beads, discs of microplates, or any other surface suitable for conducting an immunoassay. The binding processes are well known in the art and generally consist of cross-linking covalently binding or physically adsorbing, the polymer-antibody complex is washed in preparation for the test sample. An aliquot of the sample to be tested is then added to the solid phase complex and incubated for a period of time sufficient and under suitable conditions to allow binding of any antigen present to the antibody. Following the incubation period, the antigen-antibody complex is washed and dried and incubated with a second antibody specific for a portion of the antigen. The second antibody has generally a reporter molecule associated therewith that is used to indicate the binding of the second antibody to the antigen. The amount of labelled antibody that binds, as determined by the associated reporter molecule, is proportional to the amount of antigen bound to the immobilised first antibody.

An alternative method involves immobilising the antigen in the biological sample and then exposing the immobilised antigen to specific antibody that may or may not be labelled with a reporter molecule. Depending on the amount of target and the strength of the reporter molecule signal, a bound antigen may be detectable by direct labelling with the antibody. Alternatively, a second labelled antibody, specific to the first antibody is exposed to the target-first antibody complex to form a target-first antibody-second antibody tertiary complex. The complex is detected by the signal emitted by the reporter molecule.

The association of the reporter molecule with the antigen-binding molecule may include the following: (a) direct attachment of the reporter molecule to the antigen-binding molecule; (b) indirect attachment of the reporter molecule to the antigen-binding molecule; *i.e.*, attachment of the reporter molecule to another assay reagent which subsequently binds to the antigen-binding molecule; and (c) attachment to a subsequent reaction product of the antigen-binding molecule.

- 33 -

The reporter molecule may be selected from a group including a chromogen, a catalyst, an enzyme, a fluorochrome, a chemiluminescent molecule, a lanthanide ion such as Europium (Eu^{34}), a radioisotope and a direct visual label.

In the case of a direct visual label, use may be made of a colloidal metallic or non-metallic particle, a dye particle, an enzyme or a substrate, an organic polymer, a latex particle, a liposome, or other vesicle containing a signal producing substance and the like.

A large number of enzymes suitable for use as reporter molecules is disclosed in United States Patent Specifications U.S. 4,366,241, U.S. 4,843,000, and U.S. 4,849,338. Suitable enzymes useful in the present invention include alkaline phosphatase, horseradish peroxidase, luciferase, β -galactosidase, glucose oxidase, lysozyme, malate dehydrogenase and the like. The enzymes may be used alone or in combination with a second enzyme that is in solution.

Suitable fluorochromes include, but are not limited to, fluorescein isothiocyanate (FITC), tetramethylrhodamine isothiocyanate (TRITC), R-Phycoerythrin (RPE), and Texas Red. Other exemplary fluorochromes include those discussed by Dower *et al.* (International Publication WO 93/06121). Reference also may be made to the fluorochromes described in U.S. Patents 5,573,909 (Singer *et al.*), 5,326,692 (Brinkley *et al.*). Alternatively, reference may be made to the fluorochromes described in U.S. Patent Nos. 5,227,487, 5,274,113, 5,405,975, 5,433,896, 5,442,045, 5,451,663, 5,453,517, 5,459,276, 5,516,864, 5,648,270 and 5,723,218.

In the case of an enzyme immunoassay, an enzyme is conjugated to the second antibody, generally by means of glutaraldehyde or periodates. As will be readily recognised, however, a wide variety of different conjugation techniques exist which are readily available to the skilled artisan. The substrates to be used with the specific enzymes are generally chosen for the production of, upon hydrolysis by the corresponding enzyme, a detectable colour change. Examples of suitable enzymes include those described *supra*. It is also possible to employ fluorogenic substrates, which yield a fluorescent product rather than the chromogenic substrates noted above. In all cases, the enzyme-labelled antibody is added to the first antibody-antigen complex. It is then allowed to bind, and excess reagent is washed away. A solution containing the appropriate substrate is then added to the complex of antibody-antigen-antibody. The substrate will react with the enzyme linked to the second antibody, giving a qualitative visual signal, which may be further quantitated,

- 34 -

usually spectrophotometrically, to give an indication of the amount of antigen which was present in the sample.

Alternately, fluorescent compounds, such as fluorescein, rhodamine and the lanthanide, europium (EU), may be chemically coupled to antibodies without altering their binding capacity. When activated by illumination with light of a particular wavelength, the fluorochrome-labelled antibody adsorbs the light energy, inducing a state of excitability in the molecule, followed by emission of the light at a characteristic colour visually detectable with a light microscope. The fluorescent-labelled antibody is allowed to bind to the first antibody-antigen complex. After washing off the unbound reagent, the remaining tertiary complex is then exposed to light of an appropriate wavelength. The fluorescence observed indicates the presence of the antigen of interest. Immunofluorometric assays (IFMA) are well established in the art. However, other reporter molecules, such as radioisotope, chemiluminescent or bioluminescent molecules may also be employed.

It will be well understood that other means of testing target polypeptide (*e.g.*, IGFBP) levels are available, including, for instance, those involving testing for an altered level of IGFBP binding activity to an insulin-like growth factor (IGF), or Western blot analysis of IGFBP protein levels in tissues, cells or fluids using anti-IGFBP antigen-binding molecule, or assaying the amount of antigen-binding molecule or other IGFBP binding partner which is not bound to a sample, and subtracting from the total amount of antigen-binding molecule or binding partner added.

3. Identification of target molecule modulators

The invention also provides agents that modulate the level and/or functional activity of an expression product of a gene selected from an *IGFBP* gene or a gene relating to the same regulatory or biosynthetic pathway as an *IGFBP* gene. The method includes contacting a preparation comprising target molecules selected from (1) all or part of an expression product of an *IGFBP* gene or a gene relating to the same regulatory or biosynthetic pathway as the *IGFBP* gene, or a variant or derivative of that expression product, or (2) a genetic sequence which regulates expression of the expression product mentioned in (1) and which is operably connected to a marker gene, with a test agent and detecting a change in the level and/or functional activity of all or part of that expression product or variant or derivative thereof or a change in the level of expression of the marker gene or a change in the phenotype of a cell.

- 35 -

In accordance with the present invention, agents that test positive for such interaction are useful for treating and/or modulating tumorigenesis and especially for treating or preventing a liver cancer. Assays of a suitable nature for detecting, measuring or otherwise determining modulation of tumorigenesis (e.g., such as by detecting cell proliferation) are known to persons of skill in the art. For example, tumorigenesis-modulating agents could be tested for their ability to modulate cell proliferation. Typically, for cell proliferation, cell number is determined, directly, by microscopic or electronic enumeration, or indirectly, by the use of chromogenic dyes, incorporation of radioactive precursors or measurement of metabolic activity of cellular enzymes. An exemplary cell proliferation assay comprises culturing cells in the presence or absence of a test compound, and detecting cell proliferation by, for example, measuring incorporation of tritiated thymidine or by colorimetric assay based on the metabolic breakdown of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT) (Mosman, 1983, *J. Immunol. Meth.* **65**: 55-63). Cancer or tumour markers are known for a variety of cell or tissue types. Cells or tissues expressing cancer or tumour markers may be detected using monoclonal antibodies, polyclonal antisera or other antigen-binding molecules that are immuno-interactive with these tumour markers or by using nucleic acid analysis techniques, including, for example, detecting the level or presence of tumour marker-encoding polynucleotides.

Modulators contemplated by the present invention includes agonists and antagonists of *IGFBP* gene expression include antisense molecules, ribozymes and co-suppression molecules. Agonists include molecules which increase promoter activity or interfere with negative mechanisms. Agonists of *IGFBP* include molecules which overcome any negative regulatory mechanism. Antagonists of *IGFBP* polypeptides include antibodies and inhibitor peptide fragments.

Candidate agents encompass numerous chemical classes, though typically they are organic molecules, preferably small organic compounds having a molecular weight of more than 50 and less than about 2,500 Dalton. Candidate agents comprise functional groups necessary for structural interaction with proteins, particularly hydrogen bonding, and typically include at least an amine, carbonyl, hydroxyl or carboxyl group, preferably at least two of the functional chemical groups. The candidate agents often comprise cyclical carbon or heterocyclic structures and/or aromatic or polyaromatic structures substituted with one or more of the above functional groups. Candidate agents are also found among

- 36 -

biomolecules including, but not limited to: peptides, saccharides, fatty acids, steroids, purines, pyrimidines, derivatives, structural analogues or combinations thereof.

Small (non-peptide) molecule modulators of IGFBP-1, -2 and -3 are particularly preferred. In this regard, small molecules are particularly preferred because such molecules are more readily absorbed after oral administration, have fewer potential antigenic determinants, and/or are more likely to cross the cell membrane than larger, protein-based pharmaceuticals. Small organic molecules may also have the ability to gain entry into an appropriate cell and affect the expression of a gene (*e.g.*, by interacting with the regulatory region or transcription factors involved in gene expression); or affect the activity of a gene by inhibiting or enhancing the binding of accessory molecules.

Alternatively, libraries of natural compounds in the form of bacterial, fungal, plant and animal extracts are available or readily produced. Additionally, natural or synthetically produced libraries and compounds are readily modified through conventional chemical, physical and biochemical means, and may be used to produce combinatorial libraries. Known pharmacological agents may be subjected to directed or random chemical modifications, such as acylation, alkylation, esterification, amidification, etc. to produce structural analogues.

Screening may also be directed to known pharmacologically active compounds and chemical analogues thereof.

Screening for modulatory agents according to the invention can be achieved by any suitable method. For example, the method may include contacting a cell comprising a polynucleotide corresponding to an *IGFBP* gene or a gene belonging to the same regulatory or biosynthetic pathway as an *IGFBP* gene, with an agent suspected of having said modulatory activity and screening for the modulation of the level and/or functional activity of a protein encoded by said polynucleotide, or the modulation of the level of a transcript encoded by the polynucleotide, or the modulation of the activity or expression of a downstream cellular target of said protein or said transcript. Detecting such modulation can be achieved utilising techniques including, but not restricted to, ELISA, cell-based ELISA, filter-binding ELISA, inhibition ELISA, Western blots, immunoprecipitation, slot or dot blot assays, immunostaining, RIA, scintillation proximity assays, fluorescent immunoassays using antigen-binding molecule conjugates or antigen conjugates of fluorescent substances such as fluorescein or rhodamine, Ouchterlony double diffusion analysis, immunoassays employing an avidin-biotin or a streptavidin-biotin detection

- 37 -

system, and nucleic acid detection assays including reverse transcriptase polymerase chain reaction (RT-PCR).

It will be understood that a polynucleotide from which a target molecule of interest is regulated or expressed may be naturally occurring in the cell which is the subject of testing or it may have been introduced into the host cell for the purpose of testing. Further, the naturally-occurring or introduced polynucleotide may be constitutively expressed – thereby providing a model useful in screening for agents which downregulate expression of an encoded product of the sequence wherein said downregulation can be at the nucleic acid or protein level – or may require activation – thereby providing a model useful in screening for agents that upregulate expression of an encoded product of the sequence. Further, to the extent that a polynucleotide is introduced into a cell, that polynucleotide may comprise the entire coding sequence which codes for a target protein or it may comprise a portion of that coding sequence (*e.g.* a domain such as an IGF-binding domain) or a portion that regulates expression of a product encoded by the polynucleotide (*e.g.*, a promoter). For example, the promoter that is naturally associated with the polynucleotide may be introduced into the cell that is the subject of testing. In this regard, where only the promoter is utilised, detecting modulation of the promoter activity can be achieved, for example, by operably linking the promoter to a polynucleotide encoding a suitable marker including, but not restricted to, green fluorescent protein (GFP), luciferase, β -galactosidase and catecholamine acetyl transferase (CAT). Modulation of expression may be determined by measuring the activity associated with the marker-encoding polynucleotide.

In another example, the subject of detection could be a downstream regulatory target of the target molecule, rather than target molecule itself or a marker-encoding polynucleotide operably linked to a promoter of a gene encoding a product the expression of which is regulated by the target protein.

These methods provide a mechanism for performing high throughput screening of putative modulatory agents such as proteinaceous or non-proteinaceous agents comprising synthetic, combinatorial, chemical and natural libraries. These methods will also facilitate the detection of agents which bind either the polynucleotide encoding the target molecule or which modulate the expression of an upstream molecule, which subsequently modulates the expression of the polynucleotide encoding the target molecule. Accordingly, these

- 38 -

methods provide a mechanism of detecting agents that either directly or indirectly modulate the expression and/or activity of a target molecule according to the invention.

In a series of preferred embodiments, the present invention provides assays for identifying small molecules or other compounds (*i.e.*, modulatory agents) which are capable of inducing or inhibiting the level and/or or functional activity of target molecules according to the invention. The assays may be performed *in vitro* using non-transformed cells, immortalised cell lines, or recombinant cell lines. In addition, the assays may detect the presence of increased or decreased expression of genes or production of proteins on the basis of increased or decreased mRNA expression (using, for example, nucleic acid probes corresponding or complementary to a transcript of an *IGFBP* gene or of a gene relating to the same regulatory or biosynthetic pathway as the *IGFBP* gene), increased or decreased levels of protein products (using, for example, antigen binding molecules that are immuno-interactive with a polypeptide encoded by the aforementioned genes), or increased or decreased levels of expression of a marker gene (*e.g.*, GFP, β -galactosidase or luciferase) operatively linked to a target molecule-related gene regulatory region in a recombinant construct.

Thus, for example, one may culture cells which produce a particular target molecule and add to the culture medium one or more test compounds. After allowing a sufficient period of time (*e.g.*, 6-72 hours) for the compound to induce or inhibit the level and/or functional activity of the target molecule, any change in said level from an established baseline may be detected using any of the techniques described above and well known in the art. In particularly preferred embodiments, the cells are epithelial cells. Using suitable nucleic acid probes and/or antigen-binding molecules as for example described above, detection of changes in the level and or functional activity of a target molecule, and thus identification of the compound as agonist or antagonist of the target molecule, requires only routine experimentation.

In particularly preferred embodiments, a recombinant assay is employed in which a marker gene encoding, for example, GFP, β -galactosidase or luciferase is operably linked to the 5' regulatory regions of a target molecule related gene. Such regulatory regions may be easily isolated and cloned by one of ordinary skill in the art in light of the present disclosure. The marker gene and regulatory regions are joined in-frame (or in each of the three possible reading frames) so that transcription and translation of the marker gene may proceed under the control of the regulatory elements of the target molecule related gene.

- 39 -

The recombinant construct may then be introduced into any appropriate cell type although mammalian cells are preferred, and human cells are most preferred. The transformed cells may be grown in culture and, after establishing the baseline level of expression of the marker gene, test compounds may be added to the medium. The ease of detection of the expression of the marker gene provides for a rapid, high throughput assay for the identification of agonists or antagonists of the target molecules of the invention.

Compounds identified by this method will have potential utility in modifying the expression of target molecule related genes *in vivo*. These compounds may be further tested in the animal models to identify those compounds having the most potent *in vivo* effects. In addition, as described above with respect to small molecules having target polypeptide binding activity, these molecules may serve as "lead compounds" for the further development of pharmaceuticals by, for example, subjecting the compounds to sequential modifications, molecular modelling, and other routine procedures employed in rational drug design.

In another embodiment, a method of identifying agents that inhibit IGFBP activity is provided in which a purified preparation of an IGFBP protein in the presence and absence of a candidate agent under conditions in which the IGFBP is active, and the level of IGFBP activity is measured by a suitable assay. For example, an IGFBP inhibitor can be identified by measuring the ability of a candidate agent to decrease IGFBP activity in a cell (*e.g.*, a liver cell). In this method, a cell that is capable of expressing an *IGFBP* is exposed to, or cultured in the presence and absence of, the candidate agent under conditions in which the IGFBP is active in the cell, and an activity such as tumorigenesis is detected. An agent tests positive if it inhibits any of these activities.

In yet another embodiment, random peptide libraries consisting of all possible combinations of amino acids attached to a solid phase support may be used to identify peptides that are able to bind to a target molecule or to a functional domain thereof. Identification of molecules that are able to bind to a target molecule may be accomplished by screening a peptide library with a recombinant soluble target molecule. The target molecule may be purified, recombinantly expressed or synthesised by any suitable technique. Such molecules may be conveniently prepared by a person skilled in the art using standard protocols as for example described in Sambrook, *et al.*, (1989, *supra*) in particular Sections 16 and 17; Ausubel *et al.*, (1994-1998, *supra*), in particular Chapters 10 and 16; and Coligan *et al.*, (1995-1997, *supra*), in particular Chapters 1, 5 and 6.

- 40 -

Alternatively, a target polypeptide according to the invention may be synthesised using solution synthesis or solid phase synthesis as described, for example, in Chapter 9 of Atherton and Shephard (*supra*) and in Roberge *et al* (1995, *Science* **269**: 202).

To identify and isolate the peptide/solid phase support that interacts and forms a complex with a target molecule, preferably a target polypeptide, it may be necessary to label or “tag” the target polypeptide. The target polypeptide may be conjugated to any suitable reporter molecule, including enzymes such as alkaline phosphatase and horseradish peroxidase and fluorescent reporter molecules such as fluorescein isothiocyanate (FITC), phycoerythrin (PE) and rhodamine. Conjugation of any given reporter molecule, with target polypeptide, may be performed using techniques that are routine in the art. Alternatively, target polypeptide expression vectors may be engineered to express a chimeric target polypeptide containing an epitope for which a commercially available antigen-binding molecule exists. The epitope specific antigen-binding molecule may be tagged using methods well known in the art including labelling with enzymes, fluorescent dyes or coloured or magnetic beads.

For example, the “tagged” target polypeptide conjugate is incubated with the random peptide library for 30 minutes to one hour at 22° C to allow complex formation between target polypeptide and peptide species within the library. The library is then washed to remove any unbound target polypeptide. If the target polypeptide has been conjugated to alkaline phosphatase or horseradish peroxidase the whole library is poured into a petri dish containing a substrate for either alkaline phosphatase or peroxidase, for example, 5-bromo-4-chloro-3-indoyl phosphate (BCIP) or 3,3',4,4'-diaminobenzidine (DAB), respectively. After incubating for several minutes, the peptide/solid phase-target polypeptide complex changes colour, and can be easily identified and isolated physically under a dissecting microscope with a micromanipulator. If a fluorescently tagged target polypeptide has been used, complexes may be isolated by fluorescent activated sorting. If a chimeric target polypeptide having a heterologous epitope has been used, detection of the peptide/target polypeptide complex may be accomplished by using a labelled epitope specific antigen-binding molecule. Once isolated, the identity of the peptide attached to the solid phase support may be determined by peptide sequencing.

4. Method of modulating IGFBP activity

The invention also extends to a method for modulating liver cell proliferation or tumorigenesis, comprising contacting said cell with an agent for a time and under

- 41 -

conditions sufficient to modulate the level and/or functional activity of an *IGFBP* expression product, especially an IGFBP polypeptide.

In the following embodiments, an agent is suitably used to reduce, repress or otherwise inhibit cell proliferation or tumorigenesis. In one embodiment, the agent increases the level and/or functional activity of IGFBP-3. Any suitable *IGFBP-3* inducers or stabilising/activating agents may be used in this regard and these can be identified or produced by methods for example disclosed in Section 3. In an alternate embodiment, the agent decreases the level and/or functional activity of IGFBP-1 or IGFBP-2. Suitable IGFBP-1 or IGFBP-2 inhibitors may be identified or produced by methods for example disclosed in Section 3. For example, a suitable IGFBP inhibitor may comprise oligoribonucleotide sequences, that include anti-sense RNA and DNA molecules and ribozymes that function to inhibit the translation of IGFBP protein-encoding mRNA.

Anti-sense RNA and DNA molecules act to directly block the translation of mRNA by binding to targeted mRNA and preventing protein translation. In regard to antisense DNA, oligodeoxyribonucleotides derived from the translation initiation site, *e.g.*, between -10 and +10 regions of a gene encoding a polypeptide according to the invention, are preferred. Ribozymes are enzymatic RNA molecules capable of catalysing the specific cleavage of RNA. The mechanism of ribozyme action involves sequence specific hybridisation of the ribozyme molecule to complementary target RNA, followed by an endonucleolytic cleavage. Within the scope of the invention are engineered hammerhead motif ribozyme molecules that specifically and efficiently catalyse endonucleolytic cleavage of *IGFBP*, especially *IGFBP-1* or *-2* RNA sequences. Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites, which include the following sequences, GUA, GUU and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides corresponding to the region of the target gene containing the cleavage site may be evaluated for predicted structural features such as secondary structure that may render the oligonucleotide sequence unsuitable. The suitability of candidate targets may also be evaluated by testing their accessibility to hybridisation with complementary oligonucleotides, using ribonuclease protection assays.

Both anti-sense RNA and DNA molecules and ribozymes may be prepared by any method known in the art for the synthesis of RNA molecules. These include techniques for chemically synthesising oligodeoxyribonucleotides well known in the art such as for

- 42 -

example solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by *in vitro* and *in vivo* transcription of DNA sequences encoding the antisense RNA molecule. Such DNA sequences may be incorporated into a wide variety of vectors which incorporate suitable RNA polymerase promoters such as the T7 or SP6
5 polymerase promoters. Alternatively, antisense cDNA constructs that synthesise antisense RNA constitutively or inducibly, depending on the promoter used, can be introduced stably into cell lines.

Various modifications to the DNA molecules may be introduced as a means of increasing intracellular stability and half-life. Possible modifications include but are not
10 limited to the addition of flanking sequences of ribo- or deoxy- nucleotides to the 5' and/or 3' ends of the molecule or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the oligodeoxyribonucleotide backbone.

5. *Methods of preparing an IGFBP polypeptide*

An IGFBP polypeptide or variant thereof may be prepared by any suitable
15 procedure known to those of skill in the art. For example, the polypeptides may be prepared by a procedure including the steps of (a) preparing a recombinant polynucleotide comprising a nucleotide sequence encoding a polypeptide comprising the sequence set forth in any one of SEQ ID NO: 2, 4 or 6, or variant or derivative of these, which nucleotide sequence is operably linked to transcriptional and translational regulatory
20 nucleic acid; (b) introducing the recombinant polynucleotide into a suitable host cell; (c) culturing the host cell to express recombinant polypeptide from said recombinant polynucleotide; and (d) isolating the recombinant polypeptide. Preferred nucleotide sequences include, but are not limited to the sequences set forth in SEQ ID NO: 1, 3 and 5.

The recombinant polynucleotide is preferably in the form of an expression vector
25 that may be a self-replicating extra-chromosomal vector such as a plasmid, or a vector that integrates into a host genome. The transcriptional and translational regulatory nucleic acid will generally be appropriate for the host cell used for expression. Numerous types of appropriate expression vectors and suitable regulatory sequences are known in the art for a variety of host cells. Typically, the transcriptional and translational regulatory nucleic acid
30 may include, but is not limited to, promoter sequences, leader or signal sequences, ribosomal binding sites, transcriptional start and stop sequences, translational start and termination sequences, and enhancer or activator sequences. Constitutive or inducible promoters as known in the art are contemplated by the invention. The promoters may be

- 43 -

either naturally occurring promoters, or hybrid promoters that combine elements of more than one promoter.

In a preferred embodiment, the expression vector contains a selectable marker gene to allow the selection of transformed host cells. Selection genes are well known in the art and will vary with the host cell used.

The expression vector may also include a fusion partner (typically provided by the expression vector) so that the recombinant polypeptide of the invention is expressed as a fusion polypeptide with said fusion partner. The main advantage of fusion partners is that they assist identification and/or purification of said fusion polypeptide. In order to express said fusion polypeptide, it is necessary to ligate a polynucleotide according to the invention into the expression vector so that the translational reading frames of the fusion partner and the polynucleotide coincide. Well known examples of fusion partners include, but are not limited to, glutathione-S-transferase (GST), Fc portion of human IgG, maltose binding protein (MBP) and hexahistidine (HIS₆), which are particularly useful for isolation of the fusion polypeptide by affinity chromatography. For the purposes of fusion polypeptide purification by affinity chromatography, relevant matrices for affinity chromatography are glutathione-, amylose-, and nickel- or cobalt-conjugated resins respectively. Many such matrices are available in "kit" form, such as the QIAexpress™ system (Qiagen) useful with (HIS₆) fusion partners and the Pharmacia GST purification system. In a preferred embodiment, the recombinant polynucleotide is expressed in the commercial vector pFLAG as described more fully hereinafter.

Another fusion partner well known in the art is green fluorescent protein (GFP). This fusion partner serves as a fluorescent "tag" which allows the fusion polypeptide of the invention to be identified by fluorescence microscopy or by flow cytometry. The GFP tag is useful when assessing subcellular localisation of the fusion polypeptide of the invention, or for isolating cells which express the fusion polypeptide of the invention. Flow cytometric methods such as fluorescence activated cell sorting (FACS) are particularly useful in this latter application.

Preferably, the fusion partners also have protease cleavage sites, such as for Factor X_a or Thrombin, which allow the relevant protease to partially digest the fusion polypeptide of the invention and thereby liberate the recombinant polypeptide of the invention therefrom. The liberated polypeptide can then be isolated from the fusion partner by subsequent chromatographic separation.

- 44 -

Fusion partners according to the invention also include within their scope “epitope tags”, which are usually short peptide sequences for which a specific antibody is available. Well known examples of epitope tags for which specific monoclonal antibodies are readily available include c-Myc, influenza virus, haemagglutinin and FLAG tags.

5 The step of introducing into the host cell the recombinant polynucleotide may be effected by any suitable method including transfection, and transformation, the choice of which will be dependent on the host cell employed. Such methods are well known to those of skill in the art.

10 Recombinant polypeptides of the invention may be produced by culturing a host cell transformed with an expression vector containing nucleic acid encoding a polypeptide, biologically active fragment, variant or derivative according to the invention. The conditions appropriate for protein expression will vary with the choice of expression vector and the host cell. This is easily ascertained by one skilled in the art through routine experimentation.

15 Suitable host cells for expression may be prokaryotic or eukaryotic. One preferred host cell for expression of a polypeptide according to the invention is a bacterium. The bacterium used may be *Escherichia coli*. Alternatively, the host cell may be an insect cell such as, for example, *SF9* cells that may be utilised with a baculovirus expression system.

20 The recombinant protein may be conveniently prepared by a person skilled in the art using standard protocols as for example described in Sambrook, *et al.*, 1989, in particular Sections 16 and 17; Ausubel *et al.*, (1994-1998), in particular Chapters 10 and 16; and Coligan *et al.*, (1995-1997), in particular Chapters 1, 5 and 6.

25 Alternatively, the polypeptide, fragments, variants or derivatives of the invention may be synthesised using solution synthesis or solid phase synthesis as described, for example, in Chapter 9 of Atherton and Shephard (*supra*) and in Roberge *et al* (1995).

6. *Compositions*

30 The modulatory agents described in Sections 3 and 4 (therapeutic agents) can be used as actives for modulating liver cell proliferation or tumorigenesis or for the treatment or prophylaxis of a liver cancer, particularly hepatocellular carcinoma, or related conditions as, for example, described below. These therapeutic agents can be administered to a patient either by themselves, or in pharmaceutical compositions where they are mixed with a suitable pharmaceutically acceptable carrier.

- 45 -

Accordingly, the invention also provides a composition for treatment and/or prophylaxis of a liver cancer or related condition, comprising an IGFB-3 polypeptide and optionally a pharmaceutically acceptable carrier. The invention also provides a composition for treatment and/or prophylaxis of a liver cancer or related condition, comprising an a modulatory agent that modulates the level and/or functional activity of an IGFBP including, but not limited to, IGFBP-1, IGFBP-2 and IGFBP-3, and optionally a pharmaceutically acceptable carrier.

In one embodiment, the composition comprises a modulatory agent that reduces the level and/or functional activity of IGFBP-1 and/or IGFBP-2, together with a pharmaceutically acceptable carrier. In another embodiment, the composition comprises a modulatory agent that enhances the level and/or functional activity of IGFBP-3, together with a pharmaceutically acceptable carrier.

Depending on the specific conditions being treated, therapeutic agents may be formulated and administered systemically or locally. Techniques for formulation and administration may be found in "Remington's Pharmaceutical Sciences," Mack Publishing Co., Easton, Pa., latest edition. Suitable routes may, for example, include oral, rectal, transmucosal, or intestinal administration; parenteral delivery, including intramuscular, subcutaneous, intramedullary injections, as well as intrathecal, direct intraventricular, intravenous, intraperitoneal, intranasal, or intraocular injections. For injection, the therapeutic agents of the invention may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks' solution, Ringer's solution, or physiological saline buffer. For transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art. Intra-muscular and subcutaneous injection is appropriate, for example, for administration of immunogenic compositions, vaccines and DNA vaccines.

The agents can be formulated readily using pharmaceutically acceptable carriers well known in the art into dosages suitable for oral administration. Such carriers enable the compounds of the invention to be formulated in dosage forms such as tablets, pills, capsules, liquids, gels, syrups, slurries, suspensions and the like, for oral ingestion by a patient to be treated. These carriers may be selected from sugars, starches, cellulose and its derivatives, malt, gelatine, talc, calcium sulphate, vegetable oils, synthetic oils, polyols, alginic acid, phosphate buffered solutions, emulsifiers, isotonic saline, and pyrogen-free water.

- 46 -

Pharmaceutical compositions suitable for use in the present invention include compositions wherein the active ingredients are contained in an effective amount to achieve its intended purpose. The dose of agent administered to a patient should be sufficient to effect a beneficial response in the patient over time such as a reduction in the symptoms associated with a liver cancer or related condition. The quantity of the agent(s) to be administered may depend on the subject to be treated inclusive of the age, sex, weight and general health condition thereof. In this regard, precise amounts of the agent(s) for administration will depend on the judgement of the practitioner. In determining the effective amount of the agent to be administered in the treatment or prophylaxis of the liver cancer or related condition, the physician may evaluate tissue levels of an IGFBP polypeptide, and progression of the disorder. In any event, those of skill in the art may readily determine suitable dosages of the therapeutic agents of the invention.

Pharmaceutical formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Optionally, the suspension may also contain suitable stabilisers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

Pharmaceutical preparations for oral use can be obtained by combining the active compounds with solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are, in particular, fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; cellulose preparations such as, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethyl-cellulose, sodium carboxymethylcellulose, and/or polyvinylpyrrolidone (PVP). If desired, disintegrating agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate. Such compositions may be prepared by any of the methods of pharmacy but all methods include the step of bringing into association one or more therapeutic agents as described above with the carrier which constitutes one or more necessary ingredients. In

- 47 -

general, the pharmaceutical compositions of the present invention may be manufactured in a manner that is itself known, *e.g.*, by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilising processes.

5 Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions may be used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for identification or to characterise different combinations of
10 active compound doses.

Pharmaceutical which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticiser, such as glycerol or sorbitol. The push-fit capsules can contain the active ingredients in admixture with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium
15 stearate and, optionally, stabilisers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilisers may be added.

Dosage forms of the therapeutic agents of the invention may also include injecting or implanting controlled releasing devices designed specifically for this purpose or other
20 forms of implants modified to act additionally in this fashion. Controlled release of an agent of the invention may be effected by coating the same, for example, with hydrophobic polymers including acrylic resins, waxes, higher aliphatic alcohols, polylactic and polyglycolic acids and certain cellulose derivatives such as hydroxypropylmethyl cellulose. In addition, controlled release may be effected by using other polymer matrices, liposomes
25 and/or microspheres.

Therapeutic agents of the invention may be provided as salts with pharmaceutically compatible counterions. Pharmaceutically compatible salts may be formed with many acids, including but not limited to hydrochloric, sulphuric, acetic, lactic, tartaric, malic, succinic, *etc.* Salts tend to be more soluble in aqueous or other protonic
30 solvents that are the corresponding free base forms.

For any compound used in the method of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. For example, a dose can be formulated in animal models to achieve a circulating concentration range that includes

- 48 -

the IC₅₀ as determined in cell culture (*e.g.*, the concentration of a test agent, which achieves a half-maximal inhibition or enhancement of IGFBP activity). Such information can be used to more accurately determine useful doses in humans.

Toxicity and therapeutic efficacy of such therapeutic agents can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, *e.g.*, for determining the LD₅₀ (the dose lethal to 50% of the population) and the ED₅₀ (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD₅₀/ED₅₀. Compounds that exhibit large therapeutic indices are preferred. The data obtained from these cell culture assays and animal studies can be used in formulating a range of dosage for use in human. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED₅₀ with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilised. The exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition. (See for example Fingl *et al.*, 1975, in "The Pharmacological Basis of Therapeutics", Ch. 1 p1).

Dosage amount and interval may be adjusted individually to provide plasma levels of the active agent which are sufficient to maintain IGFBP-inhibitory or enhancement effects. Usual patient dosages for systemic administration range from 1-2000 mg/day, commonly from 1-250 mg/day, and typically from 10-150 mg/day. Stated in terms of patient body weight, usual dosages range from 0.02-25 mg/kg/day, commonly from 0.02-3 mg/kg/day, typically from 0.2-1.5 mg/kg/day. Stated in terms of patient body surface areas, usual dosages range from 0.5-1200 mg/m²/day, commonly from 0.5-150 mg/m²/day, typically from 5-100 mg/m²/day.

Alternately, one may administer the compound in a local rather than systemic manner, for example, *via* injection of the compound directly into a tissue, which is preferably a heart muscle tissue or a liver tissue, often in a depot or sustained release formulation. Furthermore, one may administer the drug in a targeted drug delivery system, for example, in a liposome coated with tissue-specific antibody. The liposomes will be targeted to and taken up selectively by the tissue. In cases of local administration or selective uptake, the effective local concentration of the agent may not be related to plasma concentration.

- 49 -

Thus, the present invention also contemplates a method of gene therapy of a mammal. Such a method utilises a gene therapy construct which includes an isolated polynucleotide comprising a nucleotide sequence encoding an IGFBP-3 or a biologically active fragment thereof, wherein said polynucleotide is ligated into a gene therapy vector
5 which provides one or more regulatory sequences that direct expression of said polynucleotide in said mammal.

Typically, gene therapy vectors are derived from viral DNA sequences such as adenovirus, adeno-associated viruses, herpes-simplex viruses and retroviruses. Suitable gene therapy vectors currently available to the skilled person may be found, for example,
10 in Robbins *et al.*, 1998.

If “anti-sense” therapy is contemplated (*e.g.*, IGFBP-1 and/or -2), then one or more selected portions of a *IGFBP-1* and/or *IGFBP-2* nucleic acid may be oriented 3'→5' in the gene therapy vector.

Administration of the gene therapy construct to said mammal, preferably a
15 human, may include delivery *via* direct oral intake, systemic injection, or delivery to selected tissue(s) or cells, or indirectly *via* delivery to cells isolated from the mammal or a compatible donor. An example of the latter approach would be stem-cell therapy, wherein isolated stem cells having potential for growth and differentiation are transfected with the vector comprising an *IGFBP* nucleic acid. The stem-cells are cultured for a period and then
20 transferred to the mammal being treated.

Delivery of said gene therapy construct to cells or tissues of said mammal or said compatible donor may be facilitated by microprojectile bombardment, liposome mediated transfection (*e.g.*, lipofectin or lipofectamine), electroporation, calcium phosphate or DEAE-dextran-mediated transfection, for example. A discussion of suitable delivery
25 methods may be found in Chapter 9 of Ausubel *et al.*, (1994-1998, *supra*). For example, a nucleic acid encoding IGFBP-3 may be introduced into a cell to enhance the ability of that cell to differentiate or to decrease proliferation. Alternatively, this may be achieved by introducing into a cell *IGFBP-1* and/or *IGFBP-2* antisense sequences such as 3'→5' oligonucleotides.

30 In an alternate embodiment, a polynucleotide encoding a modulatory agent of the invention may be used as a therapeutic or prophylactic composition in the form of a “naked DNA” composition as is known in the art. For example, an expression vector comprising said polynucleotide operably linked to a regulatory polynucleotide (*e.g.* a promoter,

- 50 -

transcriptional terminator, enhancer *etc*) may be introduced into an animal, preferably a mammal, where it causes production of a modulatory agent *in vivo*, preferably in a heart muscle tissue or a liver tissue.

The step of introducing the expression vector into a target cell or tissue will differ
5 depending on the intended use and species, and can involve one or more of non-viral and viral vectors, cationic liposomes, retroviruses, and adenoviruses such as, for example, described in Mulligan, R.C., (1993). Such methods can include, for example:

A. Local application of the expression vector by injection (Wolff *et al.*, 1990), surgical
implantation, instillation or any other means. This method can also be used in
10 combination with local application by injection, surgical implantation, instillation or any other means, of cells responsive to the protein encoded by the expression vector so as to increase the effectiveness of that treatment. This method can also be used in combination with local application by injection, surgical implantation, instillation or any other means, of another factor or factors required for the activity of said protein.

15 B. General systemic delivery by injection of DNA, (Calabretta *et al.*, 1993), or RNA, alone or in combination with liposomes (Zhu *et al.*, 1993), viral capsids or nanoparticles (Bertling *et al.*, 1991) or any other mediator of delivery. Improved targeting might be achieved by linking the polynucleotide/expression vector to a targeting molecule (the so-called "magic bullet" approach employing, for example, an
20 antigen-binding molecule), or by local application by injection, surgical implantation or any other means, of another factor or factors required for the activity of the protein encoded by said expression vector, or of cells responsive to said protein.

C. Injection or implantation or delivery by any means, of cells that have been modified *ex vivo* by transfection (for example, in the presence of calcium phosphate: Chen *et al.*,
25 1987, or of cationic lipids and polyamines: Rose *et al.*, 1991), infection, injection, electroporation (Shigekawa *et al.*, 1988) or any other way so as to increase the expression of said polynucleotide in those cells. The modification can be mediated by plasmid, bacteriophage, cosmid, viral (such as adenoviral or retroviral; Mulligan, 1993; Miller, 1992; Salmons *et al.*, 1993) or other vectors, or other agents of modification
30 such as liposomes (Zhu *et al.*, 1993), viral capsids or nanoparticles (Bertling *et al.*, 1991), or any other mediator of modification. The use of cells as a delivery vehicle for genes or gene products has been described by Barr *et al.*, 1991 and by Dhawan *et al.*,

- 51 -

1991. Treated cells can be delivered in combination with any nutrient, growth factor, matrix or other agent that will promote their survival in the treated subject.

5 In order that the invention may be readily understood and put into practical effect, particular preferred embodiments will now be described by way of the following non-limiting examples.

EXAMPLES

EXAMPLE 1

Reduction of IGFBP-3 expression in human hepatocellular carcinoma

Methods

5 Collection of human HCC and adjacent non-tumour liver specimens

Tissue samples were obtained intra-operatively from tumours and adjacent non-tumour livers during liver resection for HCC in 28 patients treated at the Singapore General Hospital. The samples were snap frozen in liquid nitrogen and stored at -80° C until analysis. A similar set of samples was fixed in 10% formalin and paraffin embedded.

10 The diagnosis of HCC was confirmed histologically in all cases. Prior written informed consent was obtained from all patients and the study received ethics board approval at both institutions.

Immunolocalisation of IGFBP-3

Formalin- and paraffin-embedded sections were used for IGFBP-3, IGF-II, and
15 IGF-I immunolocalisation. This was performed using rabbit anti-human IGFBP-3 (1:500) (Upstate Biotechnology, Lake Placid, NY), rabbit anti-human IGF-I (3 µg/mL) (Santa Cruz Biotechnology, Santa Cruz, CA) and rabbit anti-human IGF-II (3 µg/mL) (Austral Biologicals, San Ramon, CA) antibodies as described (14). Non-specific staining was evaluated for each specimen using either a similar concentration of IgG or by absorbing the
20 primary with appropriate specific immunogen. The slides were evaluated and intensity of the staining was scored. Specific staining was semi-quantitated by assigning a score of 0 to +++ based on increasing green fluorescence intensity. The results shown in TABLE 1 represent the average score obtained from twice staining.

Cell culture

25 Human hepatoma HepG2 cells were obtained from ATCC and maintained as monolayer cultures in Modified Eagle's Media (MEM) (Gibco, Grand Island, NY) supplemented with 10% foetal calf serum (FCS) (Gibco, Grand Island, NY) growth medium. For proliferation study, confluent cultures of HepG2 cells were trypsinised and plated at 2.0×10^4 cells in 24-well plates with growth medium. After 48h, the cell
30 monolayers were rinsed twice with MEM phenol-red serum free (PSF) media and incubated further in PSF medium for 24h. After 24h, various concentrations of human

- 53 -

recombinant IGFBP-3 (hrIGFBP-3) (Celtrix, CA), human recombinant IGF-I (hrIGF-I, GroPep, Adelaide, Australia) or 25 ng/mL hrIGF-I in conjunction with various concentrations of hrIGFBP-3 were added in triplicate in SPF media for 48 h. Cell number was determined as previously described (14).

5 To determine the effect of exogenous IGF-II and autocrine IGF-II production on HepG2 cell proliferation, HepG2 cells were plated and grown as described above and the cells subsequently treated with rabbit pre-immune serum (1:800), 25 ng/mL hrIGF-II (GroPep, Adelaide, Australia), rabbit anti-human IGF-II antibody (2 µg/mL) or with combination for 48 h. Cell number was determined as described above.

10 Western Blotting

 To determine changes in the expression of IGFBP-3 and IGF-IR, snap-frozen HCC tumours and normal adjacent liver tissues were thawed and homogenised in lysis buffer (1 mM CaCl₂, 1 mM MgCl₂, 1% NP-40, 1 µg/mL leupeptin, 1 µg/mL aprotinin, 1 µM PMSF, and 100 µM NaVO₄). Proteins were subjected to Western blot analysis as
15 described (15). Blots were incubated with 1:5000 anti-human IGFBP-3 (Upstate Biotechnology), 1 µg/mL anti-human IGF-IR (Santa Cruz, CA), 0.5 µg/mL anti α-tubulin (Santa Cruz, CA) antibodies and 1:7500 horseradish peroxidase-conjugated donkey anti-mouse or anti-rabbit secondary antibody (Pierce, Rockford, Illinois). The blots were then visualised with a chemiluminescent detection system as described by the manufacturer
20 (ECL, Amersham).

Results

 In the adult human, the liver is the main source of circulating IGF-I and IGFBP-3 (12, 13). Since local expression of IGF-I is important in autocrine and paracrine stimulation and the IGFBP-3 appears to decrease the mitogenic activity of free IGF-I,
25 presumably by competing with type I IGF receptors for the ligand, the IGFBP-3 levels in the HCC samples were determined. Total tissue lysates from both HCC cancers and adjacent morphologically normal liver cells were subjected to Western blot analysis. The resulting blots were incubated with IGFBP-3 antibody. Figure 1 shows that IGFBP-3 protein (a complex spanning 38-42 kDa) was either absent or low in HCC compared to
30 adjacent normal tissues. Unlike IGFBP-3 detected in the serum, which comprises two forms (42 and 38 kDa bands) of equal intensity, the 38 kDa IGFBP-3 was the predominant form in all normal liver tissue. This band was undetectable in 28.5% (8 of 28) of HCC

- 54 -

samples examined (TABLE 1). Although the remaining tumours (20 of 28; 71.5%) expressed IGFBP-3, the level was significantly lower than that observed in normal adjacent tissue ($p < 0.01$). Densitometric scanning showed IGFBP-3 levels to be 4 to 100-fold higher in normal liver tissue than in HCC. Subsequent blotting with anti- α -tubulin antibody showed relatively equal amounts of total protein loaded per lane (Figure 1). Figure 1C also demonstrated that proteolysis of IGFBP-3 was unlikely to be responsible for the low levels or absence of IGFBP-3 from HCC samples because low molecular weight IGFBP-3 fragments were not clearly detected on Western blot analysis of the tissue samples. The IGFBP-3 fragments, however, were detected in serum samples.

To confirm the above observation, immunohistochemical staining of HCC and normal adjacent liver tissues was performed using anti-human IGFBP-3 antibody which recognised both intact and fragmented IGFBP-3 protein. IGFBP-3 protein was clearly localised to almost all cells in the adjacent normal tissue (Figure 2A). However, this signal was absent (Figure 2B) in HCC, indicating that IGFBP-3 expression was either inactivated or decreased in these neoplastic cells.

To determine if HCC also expressed IGF-I and IGF-II, HCC and adjacent normal liver tissues were stained with anti-IGF-I and anti-IGF-II antibodies, respectively. Although normal adjacent tissues stained positively for IGF-I (Figure 3A), the expression was significantly decreased in HCC (Figure 3B). Approximately 64% (18/28) and 36% (10/28) of HCC samples examined had moderate and low IGF-I expression, respectively (TABLE 1) which is consistent with a previous report (28). IGF-II staining (Figures 3C and 3D), in contrast, was approximately equal (68%; 19 of 28) or more intense (32%; 9 of 28) in HCC compared to normal liver tissue (TABLE 1). These observations are in agreement with previous reports demonstrating over-expression of IGF-II in HCC (9-11). Our data suggests that IGF autocrine and paracrine loops exist in HCC.

Since IGFs stimulate growth responses in liver cells by binding to the IGF-I receptor (IGF-IR), IGF-IR levels in HCC and normal adjacent tissue samples were examined. Approximate 39% of HCC tumours had significantly lower IGF-IR levels than non-neoplastic adjacent liver tissues ($p < 0.01$) (Figure 4, TABLE 1).

To test the hypothesis that IGF-I and IGF-II play a role in mediating tumour cell growth, HepG2 cells were treated with various concentrations of human recombinant IGF-I or IGF-II for 48 h. Figure 5A shows dose-dependent growth stimulation by IGF-I. Similar

- 55 -

effects were observed when HepG2 cells were treated with IGF-II (data not shown). Four-fold increase in cell number was observed at the dose of 25 ng/mL IGF-I ($p < 0.01$).

As IGFBP-3 protein was undetectable or lost in the majority of HCC samples (Figure 1) and IGFBP-3 has been shown to inhibit cancer cells in an IGF-independent pathway (14), the inventors attempted to demonstrate the antiproliferative action of IGFBP-3 on HepG2 cells. Approximate 40% and 52% inhibition of basal proliferation by 250 ng/mL and 500 ng/mL hrIGFBP-3, respectively ($p < 0.01$) was obtained (Figure 5B). Concentrations of rhIGFBP-3 up to 1 μ g/mL did not further inhibit proliferation.

To determine if the proliferative action of IGF-I can be attenuated in the presence of IGFBP-3, HepG2 cells were treated with hrIGFBP-3 in the presence and absence of IGF-I for 48 h. As shown in Figure 5C, treatment of HepG2 with 25 ng/mL IGF-I for 48h resulted in a 2.8-fold increase in cell number ($p < 0.01$). IGF-induced HepG2 proliferation was significantly attenuated ($p < 0.01$) in the presence of 250 ng/mL IGFBP-3 and completely abolished at the concentration of 500 ng/mL. This result suggests that IGFBP-3 attenuated IGF-I-induced HepG2 proliferation by reducing IGF-I bioavailability.

To test the hypothesis that autocrine production of IGF-II by liver cancer cells plays a role in mediating liver cancer cell growth, HepG2 cells, which has been shown to secrete IGF-II (8), were treated with anti-IGF-II, IGF-II or with a combination for 48 h. Figure 5D shows a 3-fold induction of basal proliferation by 25 ng/mL IGF-II ($p < 0.01$). This induction was significantly attenuated by IGF-II antibody ($p < 0.01$), whereas pre-immune serum was not significantly affected. IGF-II antibody alone caused a 35% reduction in cell number ($p < 0.05$). These results suggest that the rapid proliferation of liver cancer cells *in vivo* and *in vitro* may at least, in part, be a consequence of autocrine stimulation mediated by IGF-II expression.

EXAMPLE 2

Insulin-like growth factor binding protein levels in serum of hepatocellular carcinoma patients: Potential Markers for HCC

Methods

Serum collection from human HCC, prostate cancer, BPH and healthy patients.

Prior written informed consent was obtained from all patients and the study received ethics board approval at both institutions. HCC serum samples (n=116) were

- 56 -

obtained from patients from intra-venous access line during surgery (n=17) and from venous blood samples obtained from patients during index outpatient visit and on every monthly outpatient visit (n=99) at the Singapore General Hospital. The diagnosis of HCC was confirmed histologically in all cases. Serum from non-HCC (n=45), prostate cancer
5 (n=29) and benign prostate hyperplasia (n=36) were obtained from venous blood samples obtained from patients during outpatient visit at Urology Centre, Singapore General Hospital. The serum was separated from blood samples and were snap frozen in liquid nitrogen and stored at -80° C until analysis.

Western blot analysis

10 To determine changes in the expression of IGFBP-3, IGFBP-1, IGFBP-2 and alpha fetoprotein, 3.5 µL serum were subjected to Western blot analysis as described (14). Blots were incubated with rabbit anti-human IGFBP-3 (1:5000), rabbit anti-bovine IGFBP-2 (1:2500), rabbit anti-human IGFBP-1 (1:2000) and rabbit anti-human-alpha
15 fetoprotein (1:200) antibodies and 1:7500 horseradish peroxidase-conjugated donkey anti-mouse or anti-rabbit secondary antibody (Pierce, Rockford, Illinois). The blots were then visualised with a chemiluminescent detection system as described by the manufacturer (ECL, Amersham). IGFBP-1, IGFBP-2 and IGFBP-3 were purchased from (Upstate Biotechnology).

Results

20 In the adult human, the liver is the main source of circulating, IGF-I, IGFBP-1 and IGFBP-3 (12, 13). Since local expression of IGF-I is important in autocrine and paracrine stimulation and the IGFBPs appear to regulate the mitogenic activity of free IGF-I, the IGFBP-3, IGFBP-2 and IGFBP-1 levels in the HCC serum samples were determined. Serum from HCC patients were subjected to Western blot analysis. The
25 serums derived from prostate cancer, BPH and healthy patients served as controls.

While IGFBP-3 detected in the serum comprises two forms, 42 and 38 kDa, these forms were undetectable in 70% (81 of 116) of HCC serum samples examined. Although the remaining HCC serum samples had detectable IGFBP-3, the levels were significantly lower than that observed in control serum samples (Figure 6A-6P). Unlike IGFBP-3,
30 IGFBP-2 and IGFBP-1 levels were low in control serum samples. Both IGFBP-2 and IGFBP-1 levels were increased in HCC serum but not others. IGFBP-3 levels were inversely correlated with HCC while IGFBP-1 and IGFBP-2 levels were positively

- 57 -

correlated with HCC. The elevation of IGFBP-1 was more closely associated with HCC conditioned than IGFBP-2. Since AFP levels have been shown to be useful in detecting hepatocellular carcinoma (HCC), the AFP levels were determined. As shown for example in Figure 6E, approximately 20% of HCC serum samples did not show any increase in
5 AFP. Thus, the serum AFP was not always correlated with the presence of HCC.

The disclosure of every patent, patent application, and publication cited herein is hereby incorporated herein by reference in its entirety.

The citation of any reference herein should not be construed as an admission that
10 such reference is available as "Prior Art" to the instant application

Throughout the specification the aim has been to describe the preferred embodiments of the invention without limiting the invention to any one embodiment or specific collection of features. Those of skill in the art will therefore appreciate that, in light of the instant disclosure, various modifications and changes can be made in the
15 particular embodiments exemplified without departing from the scope of the present invention. All such modifications and changes are intended to be included within the scope of the appended claims.

TABLES

TABLE 1

Expression of IGFs, IGFBP-3 and IGF-IR in normal adjacent liver tissues and HCC Tumours

Genes Examined	Normal adjacent Liver Tissues (n - 28)	HCC Tumours (n = 28)
Intense staining of IGF-I	+++ (100%)	++ 18/28 (64%) + 10/28 (36%)
Intense staining of IGF-II	+++ (100%)	+++ 19/28 (68%) ++++ 9/28 (32%)
Expression levels of IGFBP-3 levels: (compared with normal and adjacent tissue of the same patient)		
• Normal	28/28 (100%)	0/28 (0%)
• Low	0/28 (0%)	8/28 (28.5%)
• Undetectable	0/28 (0%)	20/28 (71.5%)
IGF-IR expression levels: (compared with normal and adjacent tissue of the same patient)		
• Normal	28/28 (100%)	17/28 (61%)
• Low	0/28 (0%)	11/28 (39%)

BIBLIOGRAPHY

1. Okuda, K., Ohtsuki, T., Obata, H., Tomimatsu, M., Okazaki, N., Hasegawa, H., Nakajima, Y., and Ohnishi, K. Natural history of hepatocellular carcinoma and prognosis in relation to treatment. Study of 850 patients. *Cancer*, 56: 918-928, 1985.
2. Lai, E. C., Fan, S. T., Lo, C. M., Chu, K. M., Liu, C. L., and Wong, J. Hepatic resection for hepatocellular carcinoma. An audit of 343 patients. *Ann.Surg.*, 221: 291-298, 1995.
3. Takenaka, K., Kawahara, N., Yamamoto, K., Kajiyama, K., Maeda, T., Itasaka, H., Shirabe, K., Nishizaki, T., Yanaga, K., and Sugimachi, K. Results of 280 liver resections for hepatocellular carcinoma. *Arch.Surg.*, 131: 71-76, 1996.
4. Huguet, C., Stipa, F., and Gavelli, A. Primary hepatocellular cancer: Western experience. *In* L.Blumgart (ed.), *Surgery of the liver and biliary tract*, pp. 1365-1369. Churchill Livingstone: London, 2000.
5. Lai, E. and Wong, J. Hepatocellular carcinoma: the Asian experience. *In* L. Blumgart (ed.), *Surgery of the Liver and the biliary tract*, pp. 1349-1363. London: Churchill Livingstone, 1994.
6. Chan, E., Chow PKH, Tai, B. C., Martin D, and Soo KC Neoadjuvant and adjuvant thereapy for operable hepatocellular carcinoma. *Cochrane Database System Rev.*, 2: 2000.
7. Scharf, J. G., Schmidt-Sandte, W., Pahernik, S. A., Ramadori, G., Braulke, T., and Hartmann, H. Characterization of the insulin-like growth factor axis in a human hepatoma cell line (PLC). *Carcinogenesis*, 19: 2121-2128, 1998.
8. Tsai, T. F., Yauk, Y. K., Chou, C. K., Ting, L. P., Chang, C., Hu, C. P., Han, S. H., and Su, T. S. Evidence of autocrine regulation in human hepatoma cell lines. *Biochem.Biophys.Res.Comm.*, 153: 39-45, 1988.
9. Su, T. S., Liu, W. Y., Han, S. H., Jansen, M., Yang-Fen, T. L., P'eng, F. K., and Chou, C. K. Transcripts of the insulin-like growth factors I and II in human hepatoma. *Cancer Res.*, 49: 1773-1777, 1989.

- 60 -

10. Ng, I. O., Lee, J. M., Srivastava, G., and Ng, M. Expression of insulin-like growth factor II mRNA in hepatocellular carcinoma. *J.Gastroenterol.Hepatol.*, 13: 152-157, 1998.
11. Cariani, E., Lasserre, C., Seurin, D., Hamelin, B., Kemeny, F., Franco, D., Czech, M. P., Ullrich, A., and Brechot, C. Differential expression of insulin-like growth factor II mRNA in human primary liver cancers, benign liver tumors, and liver cirrhosis. *Cancer Res.*, 48: 6844-6849, 1988.
12. Khandwala, H. M., McCutcheon, I. E., Flyvbjerg, A., and Friend, K. E. The effects of insulin-like growth factors on tumorigenesis and neoplastic growth [In Process Citation]. *Endocr.Rev.*, 21: 215-244, 2000.
13. Clemmons, D. R. Insulin-like growth factor binding proteins and their role in controlling IGF actions. *Cytokine and Growth Factor Reviews*, 8: 45-62, 1997.
14. Huynh, H., Larsson, C., Narod, S., and Pollak, M. Tumour suppressor activity of the gene encoding mammary-derived growth inhibitor. *Cancer Res.*, 55: 2225-2231, 1995.
15. Arany, E., Afford, S., Strain, A. J., Winwood, P. J., Arthur, M. J., and Hill, D. J. Differential cellular synthesis of insulin-like growth factor binding protein-1 (IGFBP-1) and IGFBP-3 within human liver. *J.Clin.Endocrinol.Metab*, 79: 1871-1876, 1994.
16. Huynh, H., Nickerson, T., Yang, X., and Pollak, M. Regulation of insulin-like growth factor I receptor by the pure antiestrogen ICI 182,780. *Clinical Cancer Res.*, 2: 2037-2042, 1996.
17. Huynh, H., Yang, X., and Pollak, M. Estradiol and antiestrogens regulate a growth inhibitory insulin-like growth factor binding protein 3 autocrine loop in human breast cancer cells. *J.Biol.Chem.*, 271: 1016-1021, 1996.
18. Oh, Y., Mullers, H. L., Ng, L., and Rosenfeld, R. G. Transforming growth factor- β -induced cell growth inhibition in human breast cancer cells is mediated through insulin-like growth factor-binding protein-3 action. *J.Biol.Chem.*, 270: 13589-13592, 1995.

- 61 -

19. LeRoith, D., Adamo, M. L., Shemer, J., Lanau, F., Shen-Orr, Z., Yaron, A., Roberts, C. T., Clemmons, D. R., Sheikh, M. S., Shao, Z. M., and. Retinoic acid inhibits growth of breast cancer cell lines: the role of insulin-like growth factor binding proteins. *Growth Regul.*, 3: 78-80, 1993.
20. Herman, J. G., Latif, F., Weng, Y., Lerman, M. I., Zbar, B., Liu, S., Samid, D., and et al. Silencing of the VHL tumor-suppressor gene by DNA methylation in renal carcinoma. *Proc.Natl.Acad.Sci.USA*, 91: 9700-9704, 1994.
21. Huynh, H., Alpert, L., and Pollak, M. Silencing of the mammary derived growth inhibitor (MDGI) gene in breast neoplasms is associated with epigenetic changes. *Cancer Res.*, 56: 4865-4870, 1996.
22. Wong, I. H., Johnson, P. J., Lai, P. B., Lau, W. Y., and Lo, Y. M. Tumor-derived epigenetic changes in the plasma and serum of liver cancer patients. Implications for cancer detection and monitoring. *Ann N.Y.Acad Sci.*, 906: 102-105, 2000.
23. Kamb, A., Gruis, N. A., Weaver-Feldhaus, J., Liu, Q., Harshman, K., Tavitgian, S. V., Stockert, E., Day, R. S., III, Johnson, B. E., and Skolnick, M. H. A cell cycle regulator potentially involved in genesis of many tumor types. *Science*, 264: 436-440, 1994.
24. Montesano, R., Hainaut, P., and Wild, C. P. Hepatocellular carcinoma: from gene to public health. *J.Natl.Cancer Inst.*, 89: 1844-1851, 1997.
25. Chen, C. J., Yu, M. W., and Liaw, Y. F. Epidemiological characteristics and risk factors of hepatocellular carcinoma. *J.Gastroenterol.Hepatol.*, 12: 5294-5308, 1997.
26. Oh, Y., Muller, H. L., Lamson, G., and Rosenfeld, R. G. Insulin-like growth factor (IGF)independent action of IGF-binding protein-3 in Hs578T human breast cancer cells. *J.Biol.Chem.*, 268: 14964-14971, 1993.
27. Oh, Y., Muller, H. L., Pham, H., and Rosenfeld, R. G. Demonstration of receptors for insulin-like growth factor binding protein-3 on Hs578T human breast cancer cells. *J.Biol.Chem.*, 268: 26045-26048, 1993.
28. Wakeling, A. E., Dukes, M., and Bowler, J. A potent specific pure antiestrogen with clinical potential. *Cancer Res.*, 51: 3867-3873, 1991.

- 62 -

29. Cameron, R. G., Imaida, K., Tsuda, H., and Ito, N. Promotive effects of steroids and bile acids on hepatocarcinogenesis initiated by diethylnitrosamine. *Cancer Res.*, 42: 2426-2428, 1982.
30. Coe, J. E., Ishak, K. G., and Ross, M. J. Estrogen induction of hepatocellular carcinomas in Armenian hamsters. *Hepatology*, 11: 570-577, 1990.

CLAIMS

1. A method for detecting the presence or diagnosing the risk of a liver cancer in a patient, comprising detecting in a biological sample obtained from said patient aberrant expression of a gene encoding an insulin-like growth factor binding protein (IGFBP).
2. The method of claim 1, wherein said IGFBP is selected from IGFBP-1, IGFBP-2 or IGFBP-3.
3. The method of claim 1, wherein said aberrant expression is detected by detecting a level and/or functional activity of an expression product of an *IGFBP-1* gene, which is elevated relative to a normal reference level and/or functional activity of said expression product.
4. The method of claim 1, wherein said aberrant expression is detected by detecting a level and/or functional activity of an expression product of an *IGFBP-2* gene, which is elevated relative to a normal reference level and/or functional activity of said expression product.
5. The method of claim 1, wherein said aberrant expression is detected by detecting a level and/or functional activity of an expression product of an *IGFBP-3* gene, which is reduced relative to a normal reference level and/or functional activity of said expression product.
6. The method of any one of claims 3 to 5, wherein the level and/or functional activity of said expression product in the biological sample obtained from said patient differs by at least 10% compared to the level and/or functional activity of said expression product in a biological sample obtained from normal patients or from patients who are not afflicted with said liver cancer.
7. The method of any one of claims 3 to 5, wherein said expression product is a transcript.
8. The method of any one of claims 3 to 5, wherein said expression product is a polypeptide.
9. The method of claim 1, wherein said liver cancer is liver cell carcinoma.
10. The method of claim 1, wherein said biological sample comprises a fluid selected from whole blood, plasma, lymph, saliva, mucus secretion and respiratory fluid.

- 64 -

11. The method of claim 1, wherein said biological sample comprises a circulatory fluid selected from serum or whole blood or a fractionated portion thereof.
12. The method of claim 1, wherein said biological sample comprises serum or a fractionated portion thereof.
13. A method for detecting the presence or diagnosing the risk of a liver cancer in a patient, comprising detecting in a biological sample obtained from said patient:
 - a level and/or functional activity of an expression product of an *IGFBP-1* gene, which is elevated relative to a normal reference level and/or functional activity of said *IGFBP-1* expression product; and/or
 - a level and/or functional activity of an expression product of an *IGFBP-2* gene, which is elevated relative to a normal reference level and/or functional activity of said *IGFBP-2* expression product; and/or
 - a level and/or functional activity of an expression product of an *IGFBP-3* gene, which is reduced relative to a normal reference level and/or functional activity of said *IGFBP-3* expression product.
14. A method for detecting the presence or diagnosing the risk of a liver cancer in a patient, comprising detecting in a biological sample obtained from said patient:
 - a level and/or functional activity of an expression product of an *IGFBP-1* gene, which is elevated relative to a normal reference level and/or functional activity of said *IGFBP-1* expression product; and
 - a level and/or functional activity of an expression product of an *IGFBP-2* gene, which is elevated relative to a normal reference level and/or functional activity of said *IGFBP-2* expression product.
15. A method for detecting the presence or diagnosing the risk of a liver cancer in a patient, comprising detecting in a biological sample obtained from said patient:
 - a level and/or functional activity of an expression product of an *IGFBP-1* gene, which is elevated relative to a normal reference level and/or functional activity of said *IGFBP-1* expression product; and
 - a level and/or functional activity of an expression product of an *IGFBP-3* gene, which is reduced relative to a normal reference level and/or functional activity of said *IGFBP-3* expression product.

- 65 -

16. A method for detecting the presence or diagnosing the risk of a liver cancer in a patient, comprising detecting in a biological sample obtained from said patient:
- a level and/or functional activity of an expression product of an *IGFBP-2* gene, which is elevated relative to a normal reference level and/or functional activity of said *IGFBP-2* expression product; and
 - a level and/or functional activity of an expression product of an *IGFBP-3* gene, which is reduced relative to a normal reference level and/or functional activity of said *IGFBP-3* expression product.
17. A method for detecting the presence or diagnosing the risk of a liver cancer in a patient, comprising detecting in a biological sample obtained from said patient:
- a level and/or functional activity of an expression product of an *IGFBP-1* gene, which is elevated relative to a normal reference level and/or functional activity of said *IGFBP-1* expression product; and
 - a level and/or functional activity of an expression product of an *IGFBP-2* gene, which is elevated relative to a normal reference level and/or functional activity of said *IGFBP-2* expression product; and
 - a level and/or functional activity of an expression product of an *IGFBP-3* gene, which is reduced relative to a normal reference level and/or functional activity of said *IGFBP-3* expression product.
18. A method for prognostic assessment of a liver cancer in a patient, comprising detecting in a biological sample obtained from said patient aberrant expression of at least one gene encoding an IGFBP.
19. Use of an agent in the manufacture of a medicament for treating and/or preventing a liver cancer, wherein said agent is optionally formulated with a pharmaceutically acceptable carrier and is identifiable by a screening assay comprising:
- contacting a preparation comprising at least a portion of an expression product of a gene selected from an *IGFBP* gene or a gene relating to the same regulatory or biosynthetic pathway as the *IGFBP* gene, or a variant or derivative of said expression product, or comprising a genetic sequence which regulates expression of said expression product and which is operably connected to a marker gene, with said agent; and

- 66 -

- detecting a change in the level and/or functional activity of said at least a portion of said expression product or said variant or derivative or a change in the level of expression of said marker gene.

20. Use of an agent that reduces the level and/or functional activity of an expression product of an *IGFBP-1* gene in the manufacture of a medicament for treating and/or preventing a liver cancer, wherein said agent is optionally formulated with a pharmaceutically acceptable carrier and is identifiable by a screening assay comprising:

- contacting a preparation comprising at least a portion of an expression product of an *IGFBP-1* gene, or a variant or derivative of said expression product, or comprising a genetic sequence which regulates expression of said expression product and which is operably connected to a marker gene, with said agent; and

- detecting a reduction in the level and/or functional activity of said at least a portion of said expression product or said variant or derivative or a reduction in the level and/or functional activity of an expression product of said marker gene.

21. Use of an agent that reduces the level and/or functional activity of an expression product of an *IGFBP-2* gene in the manufacture of a medicament for treating and/or preventing a liver cancer, wherein said agent is optionally formulated with a pharmaceutically acceptable carrier and is identifiable by a screening assay comprising:

- contacting a preparation comprising at least a portion of an expression product of an *IGFBP-2* gene, or a variant or derivative of said expression product, or comprising a genetic sequence which regulates expression of said expression product and which is operably connected to a marker gene, with said agent; and

- detecting a reduction in the level and/or functional activity of said at least a portion of said expression product or said variant or derivative or a reduction in the level and/or functional activity of an expression product of said marker gene.

22. Use of an agent that increases the level and/or functional activity of an expression product of an *IGFBP-3* gene in the manufacture of a medicament for treating and/or preventing a liver cancer, wherein said agent is optionally formulated with a pharmaceutically acceptable carrier and is identifiable by a screening assay comprising:

- contacting a preparation comprising at least a portion of an expression product of an *IGFBP-3* gene, or a variant or derivative of said expression product, or comprising a

- 67 -

genetic sequence which regulates expression of said expression product and which is operably connected to a marker gene, with said agent; and

– detecting an increase in the level and/or functional activity of said at least a portion of said expression product or said variant or derivative or an increase in the level and/or functional activity of an expression product of said marker gene.

23. The use of claim 20 or claim 21, wherein said agent is an antisense oligonucleotide or ribozyme that binds to, or otherwise interacts specifically with, a polynucleotide selected from an *IGFBP-1* polynucleotide, an *IGFBP-2* polynucleotide or complement of these.
24. The use of claim 20 or claim 21, wherein said agent is an antigen-binding molecule that is immuno-interactive with IGFBP-1 and/or IGFBP-2.
25. The use of claim 22, wherein said agent comprises an IGFBP-3 polypeptide or a biologically active fragment thereof, or variant or derivative of these.
26. The use of claim 22, wherein said agent comprises a polynucleotide encoding an IGFBP-3 polypeptide or a biologically active fragment thereof, or variant or derivative of these, wherein said polynucleotide is operably connected to a regulatory polynucleotide.
27. Use of a vector in the manufacture of a medicament, which is optionally formulated with a pharmaceutically acceptable carrier, for the treatment and/or prophylaxis of a liver cancer or related condition, wherein said vector comprises at least a portion of an *IGFBP-3* polynucleotide operably linked, in the sense direction, to a regulatory polynucleotide.
28. Use of a vector in the manufacture of a medicament, which is optionally formulated with a pharmaceutically acceptable carrier, for the treatment and/or prophylaxis of a liver cancer or related condition, wherein said vector comprises an isolated *IGFBP-1* polynucleotide operably linked, in the antisense direction, to a regulatory polynucleotide.
29. Use of a vector in the manufacture of a medicament, which is optionally formulated with a pharmaceutically acceptable carrier, for the treatment and/or prophylaxis of a liver cancer or related condition, wherein said vector comprises an isolated *IGFBP-2*

- 68 -

polynucleotide operably linked, in the antisense direction, to a regulatory polynucleotide.

30. A method for modulating liver cell proliferation or tumorigenesis, comprising contacting a liver cell with an agent for a time and under conditions sufficient to modulate the level and/or functional activity of an *IGFBP* expression product, wherein said agent is optionally formulated with a pharmaceutically acceptable carrier and is identifiable by a screening assay comprising:

- contacting a preparation comprising at least a portion of an expression product of a gene selected from an *IGFBP* gene or a gene relating to the same regulatory or biosynthetic pathway as the *IGFBP* gene, or a variant or derivative of said expression product, or comprising a genetic sequence which regulates expression of said expression product and which is operably connected to a marker gene, with said agent; and

- detecting a change in the level and/or functional activity of said at least a portion of said expression product or said variant or derivative or a change in the level of expression of said marker gene.

31. A method for modulating liver cell proliferation or tumorigenesis, comprising contacting a liver cell with an agent for a time and under conditions sufficient to reduce the level and/or functional activity of an *IGFBP-1* expression product, wherein said agent is optionally formulated with a pharmaceutically acceptable carrier and is identifiable by a screening assay comprising:

- contacting a preparation comprising at least a portion of an expression product of an *IGFBP-1* gene, or a variant or derivative of said expression product, or comprising a genetic sequence which regulates expression of said expression product and which is operably connected to a marker gene, with said agent; and

- detecting a reduction in the level and/or functional activity of said at least a portion of said expression product or said variant or derivative or a reduction in the level and/or functional activity of an expression product of said marker gene.

32. A method for modulating liver cell proliferation or tumorigenesis, comprising contacting a liver cell with an agent for a time and under conditions sufficient to reduce the level and/or functional activity of an *IGFBP-2* expression product, wherein said

- 69 -

agent is optionally formulated with a pharmaceutically acceptable carrier and is identifiable by a screening assay comprising:

- contacting a preparation comprising at least a portion of an expression product of an *IGFBP-2* gene, or a variant or derivative of said expression product, or comprising a genetic sequence which regulates expression of said expression product and which is operably connected to a marker gene, with said agent; and

- detecting a reduction in the level and/or functional activity of said at least a portion of said expression product or said variant or derivative or a reduction in the level and/or functional activity of an expression product of said marker gene.

33. A method for modulating liver cell proliferation or tumorigenesis, comprising contacting a liver cell with an agent for a time and under conditions sufficient to increase the level and/or functional activity of an *IGFBP-3* expression product, wherein said agent is optionally formulated with a pharmaceutically acceptable carrier and is identifiable by a screening assay comprising:

- contacting a preparation comprising at least a portion of an expression product of an *IGFBP-3* gene, or a variant or derivative of said expression product, or comprising a genetic sequence which regulates expression of said expression product and which is operably connected to a marker gene, with said agent; and

- detecting an increase in the level and/or functional activity of said at least a portion of said expression product or said variant or derivative or an increase in the level and/or functional activity of an expression product of said marker gene.

34. A method for treatment and/or prophylaxis of a liver cancer or related condition, comprising administering to a patient in need of such treatment an effective amount of an agent that modulates the level and/or functional activity of an *IGFBP* expression product.

35. A method for treatment and/or prophylaxis of a liver cancer or related condition, comprising administering to a patient in need of such treatment an effective amount of an agent that decreases the level and/or functional activity of an *IGFBP-1* expression product.

36. A method for treatment and/or prophylaxis of a liver cancer or related condition, comprising administering to a patient in need of such treatment an effective amount of

- 70 -

an agent that decreases the level and/or functional activity of an *IGFBP-2* expression product.

37. A method for treatment and/or prophylaxis of a liver cancer or related condition, comprising administering to a patient in need of such treatment an effective amount of an agent that increases the level and/or functional activity of an *IGFBP-3* expression product.





FIGURE 1

A



B




FIGURE 2




FIGURE 3




FIGURE 4




FIGURE 5

19/12/00 Serum1




FIGURE 6A

19/12/00 Serum 2




FIGURE 6B

19/12/00 Serum 3




FIGURE 6C

19/12/00 Serum 4




FIGURE 6D

19/12/00 Serum 5




FIGURE 6E

19/12/00 Serum 6




FIGURE 6F

19/12/00 Serum7




FIGURE 6G

19/12/00 Serum 8




FIGURE 6H

28/12/00 Serum1




FIGURE 6I

28/12/00 Serum 2




FIGURE 6J

28/12/00 Serum 3




FIGURE 6K

28/12/00 Serum 6





FIGURE 6L



N: Normal
T: Prostate Cancer
B: Benign Prostate Hyperplasia

FIGURE 6M




FIGURE 6N



N: Normal

T: Prostate Cancer

B: Benign Prostate Hyperplasia

FIGURE 60




FIGURE 6P

- 1 -

SEQUENCE LISTING

<110> National Cancer Centre of Singapore (all states, except U.S.)
 Singapore General Hospital Pte Ltd (all states, except U.S.)
 Huynh, The Hung (U.S. only)
 Chow, Pierce Kah Hoe (U.S. only)
 Soo, Khee Chee (U.S. only)

<120> Polynucleotides and polypeptides linked to cancer

<130> 2521505/VPA

<140> Yet to be assigned

<141> 2002-05-03

<150> U.S. 60/288,441

<151> 2001-05-03

<160> 12

<170> PatentIn version 3.1

<210> 1

<211> 1514

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (166) .. (945)

<223>

<400> 1

atcgggccacc gccatcccat ccagcgagca tctgccgccg cgccgccgcc accctcccag 60

agagcactgg ccaccgctcc accatcactt gcccagagtt tggggccaccg cccgccgcca 120

ccagcccaga gagcatcggc ccctgtctgc tgctcgcgcc tggag atg tca gag gtc 177
 Met Ser Glu Val
 1

ccc gtt gct cgc gtc tgg ctg gta ctg ctc ctg ctg act gtc cag gtc 225
 Pro Val Ala Arg Val Trp Leu Val Leu Leu Leu Leu Thr Val Gln Val
 5 10 15 20

ggc gtg aca gcc ggc gct ccg tgg cag tgc gcg ccc tgc tcc gcc gag 273
 Gly Val Thr Ala Gly Ala Pro Trp Gln Cys Ala Pro Cys Ser Ala Glu
 25 30 35

aag ctc gcg ctc tgc ccg ccg gtg tcc gcc tcg tgc tcg gag gtc acc 321
 Lys Leu Ala Leu Cys Pro Pro Val Ser Ala Ser Cys Ser Glu Val Thr
 40 45 50

cgg tcc gcc ggc tgc ggc tgt tgc ccg atg tgc gcc ctg cct ctg ggc 369
 Arg Ser Ala Gly Cys Gly Cys Cys Pro Met Cys Ala Leu Pro Leu Gly
 55 60 65

gcc gcg tgc ggc gtg gcg act gca cgc tgc gcc cgg gga ctc agt tgc 417
 Ala Ala Cys Gly Val Ala Thr Ala Arg Cys Ala Arg Gly Leu Ser Cys
 70 75 80

- 2 -

cgc gcg ctg ccg ggg gag cag caa cct ctg cac gcc ctc acc cgc ggc	465
Arg Ala Leu Pro Gly Glu Gln Gln Pro Leu His Ala Leu Thr Arg Gly	
85 90 95 100	
caa ggc gcc tgc gtg cag gag tct gac gcc tcc gct ccc cat gct gca	513
Gln Gly Ala Cys Val Gln Glu Ser Asp Ala Ser Ala Pro His Ala Ala	
105 110 115	
gag gca ggg agc cct gaa agc cca gag agc acg gag ata act gag gag	561
Glu Ala Gly Ser Pro Glu Ser Pro Glu Ser Thr Glu Ile Thr Glu Glu	
120 125 130	
gag ctc ctg gat aat ttc cat ctg atg gcc cct tct gaa gag gat cat	609
Glu Leu Leu Asp Asn Phe His Leu Met Ala Pro Ser Glu Glu Asp His	
135 140 145	
tcc atc ctt tgg gac gcc atc agt acc tat gat ggc tcg aag gct ctc	657
Ser Ile Leu Trp Asp Ala Ile Ser Thr Tyr Asp Gly Ser Lys Ala Leu	
150 155 160	
cat gtc acc aac atc aaa aaa tgg aag gag ccc tgc cga ata gaa ctc	705
His Val Thr Asn Ile Lys Lys Trp Lys Glu Pro Cys Arg Ile Glu Leu	
165 170 175 180	
tac aga gtc gta gag agt tta gcc aag gca cag gag aca tca gga gaa	753
Tyr Arg Val Val Glu Ser Leu Ala Lys Ala Gln Glu Thr Ser Gly Glu	
185 190 195	
gaa att tcc aaa ttt tac ctg cca aac tgc aac aag aat gga ttt tat	801
Glu Ile Ser Lys Phe Tyr Leu Pro Asn Cys Asn Lys Asn Gly Phe Tyr	
200 205 210	
cac agc aga cag tgt gag aca tcc atg gat gga gag gcg gga ctc tgc	849
His Ser Arg Gln Cys Glu Thr Ser Met Asp Gly Glu Ala Gly Leu Cys	
215 220 225	
tgg tgc gtc tac cct tgg aat ggg aag agg atc cct ggg tct cca gag	897
Trp Cys Val Tyr Pro Trp Asn Gly Lys Arg Ile Pro Gly Ser Pro Glu	
230 235 240	
atc agg gga gac ccc aac tgc cag ata tat ttt aat gta caa aac tga	945
Ile Arg Gly Asp Pro Asn Cys Gln Ile Tyr Phe Asn Val Gln Asn	
245 250 255	
aaccagatga aataatgttc tgtcacgtga aatattttaag tatatagtat atttatactc	1005
tagaacatgc acattttatat atatatgtat atgtatatat atatagtaac tactttttat	1065
actccataca taacttgata tagaaagctg tttattttatt cactgtaagt ttattttttc	1125
tacacagtaa aaacttgtag tatgttaata acttgtccta tgtcaatttg tatatcatga	1185
aacacttctc atcatattgt atgtaagtaa ttgcatttct gctcttccaa agctcctgcg	1245
tctgtttttta aagagcatgg aaaaatactg cctagaaaat gcaaaatgaa ataagagaga	1305
gtagtttttc agctagtttg aaggaggacg gttaacttgt atattccacc attcacattt	1365
gatgtacatg tgtagggaaa gttaaaagtg ttgattacat aatcaaagct acctgtggtg	1425
atgttgccac ctgtttaaata gtacactgga tatgttggtta aacacgtgtc tataatggaa	1485

- 3 -

acatttacaa taaatattct gcatggaaa

1514

<210> 2
 <211> 259
 <212> PRT
 <213> Homo sapiens

<400> 2

Met Ser Glu Val Pro Val Ala Arg Val Trp Leu Val Leu Leu Leu Leu
 1 5 10 15

Thr Val Gln Val Gly Val Thr Ala Gly Ala Pro Trp Gln Cys Ala Pro
 20 25 30

Cys Ser Ala Glu Lys Leu Ala Leu Cys Pro Pro Val Ser Ala Ser Cys
 35 40 45

Ser Glu Val Thr Arg Ser Ala Gly Cys Gly Cys Cys Pro Met Cys Ala
 50 55 60

Leu Pro Leu Gly Ala Ala Cys Gly Val Ala Thr Ala Arg Cys Ala Arg
 65 70 75 80

Gly Leu Ser Cys Arg Ala Leu Pro Gly Glu Gln Gln Pro Leu His Ala
 85 90 95

Leu Thr Arg Gly Gln Gly Ala Cys Val Gln Glu Ser Asp Ala Ser Ala
 100 105 110

Pro His Ala Ala Glu Ala Gly Ser Pro Glu Ser Pro Glu Ser Thr Glu
 115 120 125

Ile Thr Glu Glu Glu Leu Leu Asp Asn Phe His Leu Met Ala Pro Ser
 130 135 140

Glu Glu Asp His Ser Ile Leu Trp Asp Ala Ile Ser Thr Tyr Asp Gly
 145 150 155 160

Ser Lys Ala Leu His Val Thr Asn Ile Lys Lys Trp Lys Glu Pro Cys
 165 170 175

Arg Ile Glu Leu Tyr Arg Val Val Glu Ser Leu Ala Lys Ala Gln Glu
 180 185 190

Thr Ser Gly Glu Glu Ile Ser Lys Phe Tyr Leu Pro Asn Cys Asn Lys
 195 200 205

- 4 -

Asn Gly Phe Tyr His Ser Arg Gln Cys Glu Thr Ser Met Asp Gly Glu
 210 215 220

Ala Gly Leu Cys Trp Cys Val Tyr Pro Trp Asn Gly Lys Arg Ile Pro
 225 230 235 240

Gly Ser Pro Glu Ile Arg Gly Asp Pro Asn Cys Gln Ile Tyr Phe Asn
 245 250 255

Val Gln Asn

<210> 3
 <211> 1433
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (118) .. (1104)
 <223>

<400> 3
 attcggggcg agggaggagg aagaagcgga ggaggcggct cccgctcgca gggccgtgca 60
 cctgcccgcc cgcccgcctg ctcgctcgcc cgccgcgccg cgctgccgac cgccagc 117
 atg ctg ccg aga gtg ggc tgc ccc gcg ctg ccg ctg ccg ccg ccg 165
 Met Leu Pro Arg Val Gly Cys Pro Ala Leu Pro Leu Pro Pro Pro Pro
 1 5 10 15
 ctg ctg ccg ctg ctg ccg ctg ctg ctg ctg cta ctg ggc gcg agt ggc 213
 Leu Leu Pro Leu Leu Pro Leu Leu Leu Leu Leu Leu Gly Ala Ser Gly
 20 25 30
 ggc ggc ggc ggg gcg cgc gcg gag gtg ctg ttc cgc tgc ccg ccc tgc 261
 Gly Gly Gly Gly Ala Arg Ala Glu Val Leu Phe Arg Cys Pro Pro Cys
 35 40 45
 aca ccc gag cgc ctg gcc gcc tgc ggg ccc ccg ccg gtt gcg ccg ccc 309
 Thr Pro Glu Arg Leu Ala Ala Cys Gly Pro Pro Pro Val Ala Pro Pro
 50 55 60
 gcc gcg gtg gcc gca gtg gcc gga ggc gcc cgc atg cca tgc gcg gag 357
 Ala Ala Val Ala Ala Val Ala Gly Gly Ala Arg Met Pro Cys Ala Glu
 65 70 75 80
 ctc gtc cgg gag ccg ggc tgc ggc tgc tgc tgc gtg tgc gcc cgg ctg 405
 Leu Val Arg Glu Pro Gly Cys Gly Cys Cys Ser Val Cys Ala Arg Leu
 85 90 95
 gag ggc gag gcg tgc ggc gtc tac acc ccg cgc tgc ggc cag ggg ctg 453
 Glu Gly Glu Ala Cys Gly Val Tyr Thr Pro Arg Cys Gly Gln Gly Leu
 100 105 110
 cgc tgc tat ccc cac ccg ggc tcc gag ctg ccc ctg cag gcg ctg gtc 501

- 6 -

agggggaaga gaaattttta tttttgaacc cctgtgtccc ttttgcataa gattaaagga 1424

aggaaaagt 1433

<210> 4

<211> 328

<212> PRT

<213> Homo sapiens

<400> 4

Met Leu Pro Arg Val Gly Cys Pro Ala Leu Pro Leu Pro Pro Pro Pro
1 5 10 15

Leu Leu Pro Leu Leu Pro Leu Leu Leu Leu Leu Leu Gly Ala Ser Gly
20 25 30

Gly Gly Gly Gly Ala Arg Ala Glu Val Leu Phe Arg Cys Pro Pro Cys
35 40 45

Thr Pro Glu Arg Leu Ala Ala Cys Gly Pro Pro Pro Val Ala Pro Pro
50 55 60

Ala Ala Val Ala Ala Val Ala Gly Gly Ala Arg Met Pro Cys Ala Glu
65 70 75 80

Leu Val Arg Glu Pro Gly Cys Gly Cys Cys Ser Val Cys Ala Arg Leu
85 90 95

Glu Gly Glu Ala Cys Gly Val Tyr Thr Pro Arg Cys Gly Gln Gly Leu
100 105 110

Arg Cys Tyr Pro His Pro Gly Ser Glu Leu Pro Leu Gln Ala Leu Val
115 120 125

Met Gly Glu Gly Thr Cys Glu Lys Arg Arg Asp Ala Glu Tyr Gly Ala
130 135 140

Ser Pro Glu Gln Val Ala Asp Asn Gly Asp Asp His Ser Glu Gly Gly
145 150 155 160

Leu Val Glu Asn His Val Asp Ser Thr Met Asn Met Leu Gly Gly Gly
165 170 175

Gly Ser Ala Gly Arg Lys Pro Leu Lys Ser Gly Met Lys Glu Leu Ala
180 185 190

Val Phe Arg Glu Lys Val Thr Glu Gln His Arg Gln Met Gly Lys Gly

- 7 -

195	200	205
Gly Lys His His Leu Gly Leu Glu Glu Pro Lys Lys Leu Arg Pro Pro 210 215 220		
Pro Ala Arg Thr Pro Cys Gln Gln Glu Leu Asp Gln Val Leu Glu Arg 225 230 235 240		
Ile Ser Thr Met Arg Leu Pro Asp Glu Arg Gly Pro Leu Glu His Leu 245 250 255		
Tyr Ser Leu His Ile Pro Asn Cys Asp Lys His Gly Leu Tyr Asn Leu 260 265 270		
Lys Gln Cys Lys Met Ser Leu Asn Gly Gln Arg Gly Glu Cys Trp Cys 275 280 285		
Val Asn Pro Asn Thr Gly Lys Leu Ile Gln Gly Ala Pro Thr Ile Arg 290 295 300		
Gly Asp Pro Glu Cys His Leu Phe Tyr Asn Glu Gln Gln Glu Ala Cys 305 310 315 320		
Gly Val His Thr Gln Arg Met Gln 325		

<210> 5
 <211> 2497
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (108) .. (983)
 <223>

<400> 5
 ggacagaggg ccgcttcttg cctggattcc acagcttcgc gccgtgtact gtcgccccat 60
 ccctgcgcgc ccagcctgcc aagcagcgtg ccccggttgc aggcgtc atg cag cgg 116
 Met Gln Arg
 1
 gcg cga ccc acg ctc tgg gcc gct gcg ctg act ctg ctg gtg ctg ctc 164
 Ala Arg Pro Thr Leu Trp Ala Ala Ala Leu Thr Leu Leu Val Leu Leu
 5 10 15
 cgc ggg ccg ccg gtg gcg cgg gct ggc gcg agc tcg ggg ggc ttg ggt 212
 Arg Gly Pro Pro Val Ala Arg Ala Gly Ala Ser Ser Gly Gly Leu Gly
 20 25 30 35
 ccc gtg gtg cgc tgc gag ccg tgc gac gcg cgt gca ctg gcc cag tgc 260

- 8 -

Pro	Val	Val	Arg	Cys 40	Glu	Pro	Cys	Asp	Ala 45	Arg	Ala	Leu	Ala	Gln 50	Cys	
gcg	cct	ccg	ccc	gcc	gtg	tgc	gcg	gag	ctg	gtg	cgc	gag	ccg	ggc	tgc	308
Ala	Pro	Pro	Pro	Ala	Val	Cys	Ala	Glu	Leu	Val	Arg	Glu	Pro	Gly	Cys	
			55					60					65			
ggc	tgc	tgc	ctg	acg	tgc	gca	ctg	agc	gag	ggc	cag	ccg	tgc	ggc	atc	356
Gly	Cys	Cys	Leu	Thr	Cys	Ala	Leu	Ser	Glu	Gly	Gln	Pro	Cys	Gly	Ile	
		70					75					80				
tac	acc	gag	cgc	tgt	ggc	tcc	ggc	ctt	cgc	tgc	cag	ccg	tcg	ccc	gac	404
Tyr	Thr	Glu	Arg	Cys	Gly	Ser	Gly	Leu	Arg	Cys	Gln	Pro	Ser	Pro	Asp	
	85					90					95					
gag	gcg	cga	ccg	ctg	cag	gcg	ctg	ctg	gac	ggc	cgc	ggg	ctc	tgc	gtc	452
Glu	Ala	Arg	Pro	Leu	Gln	Ala	Leu	Leu	Asp	Gly	Arg	Gly	Leu	Cys	Val	
100					105					110					115	
aac	gct	agt	gcc	gtc	agc	cgc	ctg	cgc	gcc	tac	ctg	ctg	cca	gcg	ccg	500
Asn	Ala	Ser	Ala	Val	Ser	Arg	Leu	Arg	Ala	Tyr	Leu	Leu	Pro	Ala	Pro	
				120					125					130		
cca	gct	cca	gga	aat	gct	agt	gag	tcg	gag	gaa	gac	cgc	agc	gcc	ggc	548
Pro	Ala	Pro	Gly	Asn	Ala	Ser	Glu	Ser	Glu	Glu	Asp	Arg	Ser	Ala	Gly	
			135					140					145			
agt	gtg	gag	agc	ccg	tcc	gtc	tcc	agc	acg	cac	cgg	gtg	tct	gat	ccc	596
Ser	Val	Glu	Ser	Pro	Ser	Val	Ser	Ser	Thr	His	Arg	Val	Ser	Asp	Pro	
		150					155					160				
aag	ttc	cac	ccc	ctc	cat	tca	aag	ata	atc	atc	atc	aag	aaa	ggg	cat	644
Lys	Phe	His	Pro	Leu	His	Ser	Lys	Ile	Ile	Ile	Ile	Lys	Lys	Gly	His	
	165					170					175					
gct	aaa	gac	agc	cag	cgc	tac	aaa	gtt	gac	tac	gag	tct	cag	agc	aca	692
Ala	Lys	Asp	Ser	Gln	Arg	Tyr	Lys	Val	Asp	Tyr	Glu	Ser	Gln	Ser	Thr	
180					185					190					195	
gat	acc	cag	aac	ttc	tcc	tcc	gag	tcc	aag	cgg	gag	aca	gaa	tat	ggc	740
Asp	Thr	Gln	Asn	Phe	Ser	Ser	Glu	Ser	Lys	Arg	Glu	Thr	Glu	Tyr	Gly	
				200					205					210		
ccc	tgc	cgt	aga	gaa	atg	gaa	gac	aca	ctg	aat	cac	ctg	aag	ttc	ctc	788
Pro	Cys	Arg	Arg	Glu	Met	Glu	Asp	Thr	Leu	Asn	His	Leu	Lys	Phe	Leu	
			215					220					225			
aat	gtg	ctg	agt	ccc	agg	ggc	gta	cac	att	ccc	aac	tgt	gac	aag	aag	836
Asn	Val	Leu	Ser	Pro	Arg	Gly	Val	His	Ile	Pro	Asn	Cys	Asp	Lys	Lys	
		230					235					240				
gga	ttt	tat	aag	aaa	aag	cag	tgt	cgc	cct	tcc	aaa	ggc	agg	aag	cgg	884
Gly	Phe	Tyr	Lys	Lys	Lys	Gln	Cys	Arg	Pro	Ser	Lys	Gly	Arg	Lys	Arg	
	245					250					255					
ggc	ttc	tgc	tgg	tgt	gtg	gat	aag	tat	ggg	cag	cct	ctc	cca	ggc	tac	932
Gly	Phe	Cys	Trp	Cys	Val	Asp	Lys	Tyr	Gly	Gln	Pro	Leu	Pro	Gly	Tyr	
260					265					270					275	
acc	acc	aag	ggg	aag	gag	gac	gtg	cac	tgc	tac	agc	atg	cag	agc	aag	980
Thr	Thr	Lys	Gly	Lys	Glu	Asp	Val	His	Cys	Tyr	Ser	Met	Gln	Ser	Lys	
				280					285					290		

- 9 -

```

tag acgcctgccg caaggttaat gtggagctca aatatgcctt attttgcaca      1033
aaagactgcc aaggacatga ccagcagctg gctacagcct cgatttatat ttctgtttgt  1093
ggatgaactga ttttttttta aaccaaagtt tagaaagagg tttttgaaat gcctatgggt  1153
tctttgaatg gtaaacttga gcatcttttc actttccagt agtcagcaaa gagcagtttg  1213
aattttcttg tcgcttecta tcaaaatatt cagagactcg agcacagcac ccatacttca  1273
tgcgcccgtg gaatgctcac cacatgttgg tcgaagcggc cgaccactga ctttgtgact  1333
taggcggctg tgttgccat gtagagaaca cgcttcaccc ccactccccg tacagtgcgc  1393
acaggcttta tcgagaatag gaaaaccttt aaaccccggt catccggaca tcccaacgca  1453
tgctcctgga gctcacagcc ttctgtgggt tcatttctga aacaagggcg tggatccctc  1513
aaccaagaag aatgtttatg tcttcaagt acctgtactg cttggggact attggagaaa  1573
ataaggtgga gtcctacttg tttaaaaaat atgtatctaa gaatgttcta gggcactctg  1633
ggaacctata aaggcaggta ttteggggcc tcctcttcag gaatcttcct gaagacatgg  1693
cccagtcgaa ggcccaggat ggcttttgct gcggccccgt ggggtaggag ggacagagag  1753
acagggagag tcagcctcca cattcagagg catcacaagt aatggcacia ttcttcggat  1813
gactgcagaa aatagtgttt tgtagttcaa caactcaaga cgaagcttat ttctgaggat  1873
aagctcttta aaggcaaagc tttattttca tctctcatct tttgtcctcc ttagcacaat  1933
gtaaaaaaga atagtaatat cagaacagga aggaggaatg gcttgctggg gagcccatcc  1993
aggacactgg gagcacatag agattcaccc atgtttgttg aacttagagt cattctcatg  2053
cttttcttta taattcacac atatatgcag agaagatatg ttcttggtta cattgtatac  2113
aacatagccc caaatatagt aagatctata ctagataatc ctagatgaaa tgtagagat  2173
gctatatgat acaactgtgg ccatgactga ggaaaggagc tcacgcccag agactgggct  2233
gctctcccgg aggccaaacc caagaaggtc tggcaaagtc aggctcaggg agactctgcc  2293
ctgctgcaga cctcgggtgtg gacacacgct gcatagagct ctccttgaaa acagaggggt  2353
ctcaagacat tctgcctacc tattagcttt tctttatttt tttaactttt tggggggaaa  2413
agtatttttg agaagtttgt cttgcaatgt atttataaat agtaaataaa gtttttacca  2473
ttaaaaaaaaa aaaaaaaaaa aaaa      2497

```

<210> 6
 <211> 291
 <212> PRT
 <213> Homo sapiens

<400> 6

Met Gln Arg Ala Arg Pro Thr Leu Trp Ala Ala Ala Leu Thr Leu Leu

1					5					10					15				
Val	Leu	Leu	Arg	Gly	Pro	Pro	Val	Ala	Arg	Ala	Gly	Ala	Ser	Ser	Gly				
			20							25				30					
Gly	Leu	Gly	Pro	Val	Val	Arg	Cys	Glu	Pro	Cys	Asp	Ala	Arg	Ala	Leu				
			35				40							45					
Ala	Gln	Cys	Ala	Pro	Pro	Pro	Ala	Val	Cys	Ala	Glu	Leu	Val	Arg	Glu				
			50				55							60					
Pro	Gly	Cys	Gly	Cys	Cys	Leu	Thr	Cys	Ala	Leu	Ser	Glu	Gly	Gln	Pro				
65							70				75								
Cys	Gly	Ile	Tyr	Thr	Glu	Arg	Cys	Gly	Ser	Gly	Leu	Arg	Cys	Gln	Pro				
						85				90				95					
Ser	Pro	Asp	Glu	Ala	Arg	Pro	Leu	Gln	Ala	Leu	Leu	Asp	Gly	Arg	Gly				
			100							105				110					
Leu	Cys	Val	Asn	Ala	Ser	Ala	Val	Ser	Arg	Leu	Arg	Ala	Tyr	Leu	Leu				
			115				120							125					
Pro	Ala	Pro	Pro	Ala	Pro	Gly	Asn	Ala	Ser	Glu	Ser	Glu	Glu	Asp	Arg				
			130				135							140					
Ser	Ala	Gly	Ser	Val	Glu	Ser	Pro	Ser	Val	Ser	Ser	Thr	His	Arg	Val				
145							150				155								
Ser	Asp	Pro	Lys	Phe	His	Pro	Leu	His	Ser	Lys	Ile	Ile	Ile	Ile	Lys				
						165				170									
Lys	Gly	His	Ala	Lys	Asp	Ser	Gln	Arg	Tyr	Lys	Val	Asp	Tyr	Glu	Ser				
			180							185				190					
Gln	Ser	Thr	Asp	Thr	Gln	Asn	Phe	Ser	Ser	Glu	Ser	Lys	Arg	Glu	Thr				
			195				200							205					
Glu	Tyr	Gly	Pro	Cys	Arg	Arg	Glu	Met	Glu	Asp	Thr	Leu	Asn	His	Leu				
			210				215							220					
Lys	Phe	Leu	Asn	Val	Leu	Ser	Pro	Arg	Gly	Val	His	Ile	Pro	Asn	Cys				
225							230				235								
Asp	Lys	Lys	Gly	Phe	Tyr	Lys	Lys	Lys	Gln	Cys	Arg	Pro	Ser	Lys	Gly				
						245				250									

- 11 -

Arg Lys Arg Gly Phe Cys Trp Cys Val Asp Lys Tyr Gly Gln Pro Leu
260 265 270

Pro Gly Tyr Thr Thr Lys Gly Lys Glu Asp Val His Cys Tyr Ser Met
275 280 285

Gln Ser Lys
290

<210> 7
<211> 6128
<212> DNA
<213> Homo sapiens

<220>
<221> mRNA
<222> (597) .. (1110)
<223> Exon 1

<220>
<221> mRNA
<222> (2657) .. (2826)
<223> Exon 2

<220>
<221> mRNA
<222> (4041) .. (4169)
<223> Exon 3

<220>
<221> mRNA
<222> (5069) .. (5769)
<223> Exon 4

<220>
<221> CDS
<222> (762) .. (1110)
<223>

<220>
<221> CDS
<222> (2657) .. (2826)
<223>

<220>
<221> CDS
<222> (4041) .. (4169)
<223>

<220>
<221> CDS

- 12 -

<222> (5069) .. (5197)

<223>

<400> 7

```

ggggagcttg gcagagccct aggatgaact gtgttttgtt ttgttttgtt ttgttgcagg      60
ttggagcctt tttcctgcag aaagagaagc aattccgggc ttttccacag ataacctgct      120
agggctcgca gagacctgga ttctaacccc atctcgctt tcttcacctg gggcattgtt      180
ttctgcattt gagaactgct ggggtgggctg gtgttaccca ggatttttaa atttttgcag      240
gagacgcttt gcaggagatc tctagggctt ttttctgtgt caattaaaga gcctggccag      300
ctggacctgg gctgtctttt tgacaaaaac aaacgtcatc cccctcccag ctgagcactt      360
gttagaactg gactttaact gagggcctga accccctaac aacgggacaa acagtatgaa      420
atcaaaaccg tttaccctcc tcccaccagc ggtttgcgta gggccttggg tgcactagca      480
aaacaaactt attttgaaca ctcagctcct agcgtgcggc gctgccaatc attaacctcc      540
tggtgcaagt ggcgcggcct gtgcccttta taagggtgcgc gctgtgtcca gcgagcatcg      600
gccaccgcca tcccatccag cgagcatctg ccgcgcgcgc gccgccaccc tcccagagag      660
cactggccac cgctccacca tcaattgccc agagtttggg ccaccgcccg ccgccaccag      720
cccagagagc atcgccctct gtctgctgct cgcgcctgga g atg tca gag gtc ccc      776
                                   Met Ser Glu Val Pro
                                   1             5

gtt gct cgc gtc tgg ctg gta ctg ctc ctg ctg act gtc cag gtc ggc      824
Val Ala Arg Val Trp Leu Val Leu Leu Leu Leu Thr Val Gln Val Gly
                        10                        15                        20

gtg aca gcc ggc gct ccg tgg cag tgc gcg ccc tgc tcc gcc gag aag      872
Val Thr Ala Gly Ala Pro Trp Gln Cys Ala Pro Cys Ser Ala Glu Lys
                        25                        30                        35

ctc gcg ctc tgc ccg ccg gtg tcc gcc tcg tgc tcg gag gtc acc cgg      920
Leu Ala Leu Cys Pro Pro Val Ser Ala Ser Cys Ser Glu Val Thr Arg
                        40                        45                        50

tcc gcc ggc tgc ggc tgt tgc ccg atg tgc gcc ctg cct ctg ggc gcc      968
Ser Ala Gly Cys Gly Cys Cys Pro Met Cys Ala Leu Pro Leu Gly Ala
                        55                        60                        65

gcg tgc ggc gtg gcg act gca cgc tgc gcc cgg gga ctc agt tgc cgc      1016
Ala Cys Gly Val Ala Thr Ala Arg Cys Ala Arg Gly Leu Ser Cys Arg
                        70                        75                        80                        85

gcg ctg ccg ggg gag cag caa cct ctg cac gcc ctc acc cgc ggc caa      1064
Ala Leu Pro Gly Glu Gln Gln Pro Leu His Ala Leu Thr Arg Gly Gln
                        90                        95                        100

ggc gcc tgc gtg cag gag tct gac gcc tcc gct ccc cat gct gca g      1110
Gly Ala Cys Val Gln Glu Ser Asp Ala Ser Ala Pro His Ala Ala
                        105                        110                        115

gtaccacagt cccgccctg ctccagcacc tctgcccgc cgcccccgcc cggcacctcc      1170

```

- 13 -

cgccccact	cccctaacta	ctgggtgggg	ctgcagccgg	gcaggggctt	gtccacaact	1230
agagcttgaa	accagagcac	gtagttgggg	aaggagcttg	ggtcacccag	tggagcccg	1290
tcattgcacg	gtcttggcag	gacgtgccct	gggagaagaa	ggaagatgtt	ccagggcaca	1350
catagcttag	tggagactca	aggaggaagc	tcccacttgg	ccctcgtagc	ccagggatct	1410
ttagagaccc	agacagcaag	agtagaggga	ggcgggtggg	ctgcctccgc	agttgggatg	1470
ccaccctggg	tccccacca	gtctccttgc	cagcacgtgg	gcagccccag	tggctggagc	1530
atccgggaag	gcctaagaag	ggaatccctc	ctgctggcct	ccccaggagg	tgtttggaat	1590
gtcctttgta	atatacgtcc	tggatacagt	atgtgcttcc	cagatgttta	cagaacataa	1650
tgtgagagtt	gaggctaaaa	gcttcacttc	cattcattgc	catagaaaaa	caatgtttga	1710
tgtgtatgtt	gccccctaag	aacatgacaa	tcaggttctt	tcctggattt	aagcaggcat	1770
ttgtttgtag	cggaagtgg	aggaaatgcc	tctttctcta	ctccagcccc	tatctcctca	1830
tctggactgc	ttgcacgtcc	tgtcctgtgc	cattccttga	ggaattagga	cttccatacc	1890
ctgatagggt	gttagagtga	gtgtgtgaga	gtcagtgacc	cttccaagga	gggaacaggg	1950
atttcctgcg	gattctctgc	ccttcagagg	aatgaaagg	gagtcgtctc	actgatcctc	2010
aaggagatgg	atttccacag	ggggtccctt	ataggccagt	gtgctgattc	tgttcatgta	2070
cctttggact	ctatctgcaa	gatcctctcg	ccatgggggt	ggaggggggtg	ttgcagagaa	2130
gtccattagg	acgtgggaag	gcagggggccc	caaaatagaa	gagggtagta	gtttgggtca	2190
ggagagctgc	attctagact	ccacttctag	atgaactgta	ccatccattc	agggatatta	2250
gtaaaatccc	cattttaagg	atggagaagc	tgaggtttcg	aggctcccta	ctaatgcggt	2310
taggaactga	actctaatac	tggcttcttt	agggttggtc	tcaacctcag	caaatgcatt	2370
tgttgctctg	ctccctgct	gctttaaaag	cagaaacgtg	gggcaccctt	gccaggccct	2430
gaaacttact	attgtcagga	acagagctgg	catgctgttg	ggacataact	cttcctggaa	2490
accattacaa	ggtcctttac	agtgagagtc	agtgaatac	aggttaggga	atgtggccat	2550
cctcatggcc	cagcctgggt	ttgagggaag	gaagtgatct	cctggggggcc	cgaaaggggc	2610
agatgctttg	ggagggtca	tgggggtctgc	aatgtttcct	ttccag ag	gca ggg	2664
				Glu Ala Gly		
agc cct gaa agc cca gag agc acg gag ata act gag gag gag ctc ctg	2712					
Ser Pro Glu Ser Pro Glu Ser Thr Glu Ile Thr Glu Glu Glu Leu Leu						
120 125 130 135						
gat aat ttc cat ctg atg gcc cct tct gaa gag gat cat tcc atc ctt	2760					
Asp Asn Phe His Leu Met Ala Pro Ser Glu Glu Asp His Ser Ile Leu						
140 145 150						
tgg gac gcc atc agt acc tat gat ggc tcg aag gct ctc cat gtc acc	2808					

- 14 -

Trp	Asp	Ala	Ile	Ser	Thr	Tyr	Asp	Gly	Ser	Lys	Ala	Leu	His	Val	Thr		
			155					160					165				
aac atc aaa aaa tgg aag gtgaggccca gcaggtgggt ggtgggaact 2856																	
Asn	Ile	Lys	Lys	Trp	Lys												
		170															
ctcctgtcag aggatgtagc ttgagtcggc tgtgacaagc agacctggcc cactccttcc 2916																	
tggtcctgat gccctggaaa gaaatgcttt tcccccaaaa cctagtggcc atttctcagt 2976																	
taaactttgt caaactgtga acacatggac tttcaggaga ggagcccact ggatgcaaag 3036																	
aaaatgtagt tccggacgat aggttacttc atgtctctga gctcctctct aaaggggtgtg 3096																	
agaacattga gttacattgt tccaaaactt agtacaaggg cctaactgct ttcaccccca 3156																	
ttgtatgatt tgctccctt taaaaaagtt gtcataaaat agacataaaa tgtactgttt 3216																	
taactatttt gaagcatagg tttctgtggc attaagtaac aataagtaat gcatgttatt 3276																	
ctgcaaccac taccaatatc taccttcaga attttcatct tcccaaactg aagctcggaa 3336																	
ctcattaaac atcaactctc cattatctta attatgtctg ttttgaattc taccacttga 3396																	
gggcgacaga gaatcaaagt cggagtgatt tgatttgaga aaataaatta cttccacaaa 3456																	
catgcctacc cagaaaaagg gaggattaaa ccattttctc aacatttttg aatttcatct 3516																	
gagaataacg ataaatggag gaaagttaat tcttaaatta tccaaatgaa tgccttaatc 3576																	
ctcacttgat ttagtttatg taaaagaacc ttgtagattg cagaggaaag ttattcaaaa 3636																	
tctagaaatg tcttctgcta aattcttaat taagagttat gccactctt ctaaccagat 3696																	
ggcgggtgcc aaaagtattc agtaggtggc aggaagagtt gggaagagtt ttgtggtggt 3756																	
ggtgaggctc ttgccagggg atgtctctgt gcaggtggtg agagagtgat tggtgaggga 3816																	
gagaatggtg ttcttggttct ttatctgaac aacttaagac agactaactg caggtgggaa 3876																	
aaagaggccc agagggagca catcacatga cacctagtcc caggtccccg tggccagatc 3936																	
gtagggccac tcctgcttct acaaaacctc ttccatcagt cattgtctag gctgataatt 3996																	
cactgtgtct tgtagatat gctaattgtca agttattctc ttag gag ccc tgc cga 4052																	
													Glu	Pro	Cys	Arg	
														175			
ata gaa ctc tac aga gtc gta gag agt tta gcc aag gca cag gag aca 4100																	
Ile	Glu	Leu	Tyr	Arg	Val	Val	Glu	Ser	Leu	Ala	Lys	Ala	Gln	Glu	Thr		
		180					185					190					
tca gga gaa gaa att tcc aaa ttt tac ctg cca aac tgc aac aag aat 4148																	
Ser	Gly	Glu	Glu	Ile	Ser	Lys	Phe	Tyr	Leu	Pro	Asn	Cys	Asn	Lys	Asn		
	195					200					205						
gga ttt tat cac agc aga cag gtaggtggcc ttgccagtgt gcgtcgtcag 4199																	
Gly	Phe	Tyr	His	Ser	Arg	Gln											
210					215												
ggtgaaaggg actactgccc tacattcctg ccaagccacg gtcattcatg tcaaagaagg 4259																	

- 15 -

tccactccaa agtagacacc agaagtgggtt gtattgagcc agatccaccc ctctgggaac	4319
ctggaacagc taggtgaaga aagcctattg aacagggtcaa tatgtcacca ccagagggtgg	4379
aaaacctgcg ctggaccagg ggccctgggg ctaagcctga gctccatcac tcagcaagcc	4439
aatagctatg gaagactttc ctagggcccc taacgcttgg atcttggggtt cctcacctat	4499
acaagagagg aagggaacaa acttaactta gaaaagcaag tatctatattg gacagctctt	4559
aacacttttt tctccttgaa ctgtctaata aaacacaaca acatttttgt ggcttgccag	4619
ataataaatg agatcaaaca gaccaagatt tcatattgag gagtgcttta ggtctcagtg	4679
aagtacaggt tctgtagatt ttattgggag aaactgagga ctaggccctg cttcacgggc	4739
aatgaacagt ggggcacaca cgagacatgt tccctctggg atgggctccc ctgacatcag	4799
gctatgaagc agacagctgt acacacactg tactgtttaa cacacatggg aagtcattat	4859
tgcacatgcc actcggtcac tgtctctatt ttatgatgag aaccaggagg gtgtgagatt	4919
tgctgcac atgggcagct ggtttcacag cggggactct tggcttggtc ctgcagtgtc	4979
cggctgcact gaaaaaaggg ccagctatgg ctctactttc cctgtcagct tgtcatctgc	5039
tgctcttgac cttgggtcttg tctttgcag tgt gag aca tcc atg gat gga gag	5092
	Cys Glu Thr Ser Met Asp Gly Glu
	220
gcg gga ctc tgc tgg tgc gtc tac cct tgg aat ggg aag agg atc cct	5140
Ala Gly Leu Cys Trp Cys Val Tyr Pro Trp Asn Gly Lys Arg Ile Pro	
225 230 235 240	
ggg tct cca gag atc agg gga gac ccc aac tgc cag atg tat ttt aat	5188
Gly Ser Pro Glu Ile Arg Gly Asp Pro Asn Cys Gln Met Tyr Phe Asn	
245 250 255	
gta caa aac tgaaaccaga tgaaataatg ttctgtcacg tgaaatatatt	5237
Val Gln Asn	
aagtatatag tatatttata ctctagaaca tgcacattta tatatatatg tatatgtata	5297
tatatatagt aactactttt tatactccat acataacttg atatagaaag ctgtttatatt	5357
attcactgta agtttatattt ttctacacag taaaaacttg tactatgtta ataacttgtc	5417
ctatgtcaat ttgtatatca tgaaacactt ctcacatcat tgtatgtaag taattgcatt	5477
tctgctcttc caaagctcct gcgtctgttt ttaaagagca tggaaaaata ctgcctagaa	5537
aatgcaaaat gaaataagag agagtagttt ttcagctagt ttgaaggagg acggttaact	5597
tgtatatattcc accattcaca tttgatgtac atgtgtaggg aaagttaaaa gtgttgatta	5657
cataatcaaa gctacctgtg gtgatgttgc cacctgttaa aatgtacact ggatatgttg	5717
ttaaacacgt gtctataatg gaaacattta caataaatat tctgcatgga aatactgtta	5777
tctctgtgga tgtttatatta gtgatattat ctttttgaaa tccctttgtc cttcacccctc	5837

- 16 -

accttcctta tgcattttgt gctttttgtga taccaagatg gtgttttggg tctgcagagc 5897
 cccttgtect aactacccta gatcttgccc cggctgcccc agggtttcca cttcccagtg 5957
 gcattggggac cttcctcctc tgcctggcca ggctgggcat tcctgtgtcc cacctctcca 6017
 ttagtcccca tcctccccag ggtcctcagc ctcttcctct tggtttggtt tgtcccttcc 6077
 gttcaactta taaataggaa cagggtttct ttcctcttaa acacaagaca g 6128

<210> 8
 <211> 259
 <212> PRT
 <213> Homo sapiens

<400> 8

Met Ser Glu Val Pro Val Ala Arg Val Trp Leu Val Leu Leu Leu Leu
 1 5 10 15

Thr Val Gln Val Gly Val Thr Ala Gly Ala Pro Trp Gln Cys Ala Pro
 20 25 30

Cys Ser Ala Glu Lys Leu Ala Leu Cys Pro Pro Val Ser Ala Ser Cys
 35 40 45

Ser Glu Val Thr Arg Ser Ala Gly Cys Gly Cys Cys Pro Met Cys Ala
 50 55 60

Leu Pro Leu Gly Ala Ala Cys Gly Val Ala Thr Ala Arg Cys Ala Arg
 65 70 75 80

Gly Leu Ser Cys Arg Ala Leu Pro Gly Glu Gln Gln Pro Leu His Ala
 85 90 95

Leu Thr Arg Gly Gln Gly Ala Cys Val Gln Glu Ser Asp Ala Ser Ala
 100 105 110

Pro His Ala Ala Glu Ala Gly Ser Pro Glu Ser Pro Glu Ser Thr Glu
 115 120 125

Ile Thr Glu Glu Glu Leu Leu Asp Asn Phe His Leu Met Ala Pro Ser
 130 135 140

Glu Glu Asp His Ser Ile Leu Trp Asp Ala Ile Ser Thr Tyr Asp Gly
 145 150 155 160

Ser Lys Ala Leu His Val Thr Asn Ile Lys Lys Trp Lys Glu Pro Cys
 165 170 175

- 17 -

Arg Ile Glu Leu Tyr Arg Val Val Glu Ser Leu Ala Lys Ala Gln Glu
180 185 190

Thr Ser Gly Glu Glu Ile Ser Lys Phe Tyr Leu Pro Asn Cys Asn Lys
195 200 205

Asn Gly Phe Tyr His Ser Arg Gln Cys Glu Thr Ser Met Asp Gly Glu
210 215 220

Ala Gly Leu Cys Trp Cys Val Tyr Pro Trp Asn Gly Lys Arg Ile Pro
225 230 235 240

Gly Ser Pro Glu Ile Arg Gly Asp Pro Asn Cys Gln Met Tyr Phe Asn
245 250 255

Val Gln Asn

<210> 9
<211> 31024
<212> DNA
<213> Homo sapiens

<220>
<221> gene
<222> (1) .. (31024)
<223>

<220>
<221> mRNA
<222> (1) .. (556)
<223> Exon 1

<220>
<221> mRNA
<222> (27148) .. (27377)
<223> Exon 2

<220>
<221> mRNA
<222> (28449) .. (28589)
<223> Exon 3

<220>
<221> mRNA
<222> (30531) .. (31024)
<223> Exon 4

<220>
<221> CDS

- 18 -

<222> (115) .. (556)
 <223>

<220>
 <221> CDS
 <222> (27148) .. (27377)
 <223>

<220>
 <221> CDS
 <222> (28449) .. (28589)
 <223>

<220>
 <221> CDS
 <222> (30531) .. (30692)
 <223>

<400> 9
 ggcgaggggag gaggaagaag cggaggaggc ggctcccgcg ctgcgagggc cgtgccacct 60
 gcccgcccgc ccgctcgctc gctcgcccgc cgcgcccgcg tgccgaccgc cagc atg 117
 Met
 1
 ctg ccg aga gtg ggc tgc ccc gcg ctg ccg ctg ccg ccg ccg ccg ctg 165
 Leu Pro Arg Val Gly Cys Pro Ala Leu Pro Leu Pro Pro Pro Pro Leu
 5 10 15
 ctg ccg ctg ctg ctg ctg cta ctg ggc gcg agt ggc ggc ggc ggc ggg 213
 Leu Pro Leu Leu Leu Leu Leu Leu Gly Ala Ser Gly Gly Gly Gly Gly
 20 25 30
 gcg cgc gcg gag gtg ctg ttc cgc tgc ccg ccc tgc aca ccc gag cgc 261
 Ala Arg Ala Glu Val Leu Phe Arg Cys Pro Pro Cys Thr Pro Glu Arg
 35 40 45
 ctg gcc gcc tgc ggg ccc ccg ccg gtt gcg ccg ccc gcc gcg gtg gcc 309
 Leu Ala Ala Cys Gly Pro Pro Pro Val Ala Pro Pro Ala Ala Val Ala
 50 55 60 65
 gca gtg gcc gga ggc gcc cgc atg cca tgc gcg gag ctc gtc cgg gag 357
 Ala Val Ala Gly Gly Ala Arg Met Pro Cys Ala Glu Leu Val Arg Glu
 70 75 80
 ccg ggc tgc ggc tgc tgc tcg gtg tgc gcc ccg ctg gag ggc gag gcg 405
 Pro Gly Cys Gly Cys Cys Ser Val Cys Ala Arg Leu Glu Gly Glu Ala
 85 90 95
 tgc ggc gtc tac acc ccg cgc tgc ggc cag ggg ctg cgc tgc tat ccc 453
 Cys Gly Val Tyr Thr Pro Arg Cys Gly Gln Gly Leu Arg Cys Tyr Pro
 100 105 110
 cac ccg ggc tcc gag ctg ccc ctg cag gcg ctg gtc atg ggc gag ggc 501
 His Pro Gly Ser Glu Leu Pro Leu Gln Ala Leu Val Met Gly Glu Gly
 115 120 125
 act tgt gag aag cgc cgg gac gcc gag tat ggc gcc agc ccg gag cag 549

- 19 -

[illegible]

- 20 -

tagccaggaa	cagatgcggc	agattttctc	ccattcactt	ttccctggag	cctgaagggg	2346
gtggggggta	gttgtcaggg	gaggccaggg	acaatggggg	gagttcatgg	gaagaatgtc	2406
actctggatt	ttctgcctgg	ttatgggact	cccttccgcc	agcttgcccc	agctgcgttc	2466
tcctccccag	tgcccagacg	ccttccgcat	ttttgtgcac	cggcctggga	agaggggatg	2526
ttaagcaggg	gtggaaggtg	gggggtgggg	ctggaagctg	agtgggtgca	ctcaaaagaa	2586
gtgaccaatg	tcaggggtgc	ctgagaaaga	gtcgggctgg	gggaggggag	cactttggga	2646
aggagcctga	gcttgcaagg	gtgcaggaga	aaaaggggtc	aaggttggag	gcttttgtgt	2706
gggagagggg	cttaggctgg	ggcagagact	ggagtcactg	gatggagagt	gctgggtccg	2766
agttgggtgg	ggcctggccc	agacggagat	cttagcactg	aatcctctgc	ctcccactcc	2826
atctcccgtc	ccgtcttctc	tcctctccga	agaatgagca	ccccccctcc	agcctgcatt	2886
cctctttgag	caggttgctt	ttgagaattt	taaacatacc	ctcttatctc	gaggcagcga	2946
acctcagcag	ttccggcttt	gctgttagct	cttttgtctt	tgttcccggg	actcaggggc	3006
tgtgttgctg	ctggcttcct	ggggatggca	aggatgcagg	ggtgggggtg	ggcacatcag	3066
tgggtggaag	gagtgggtgg	tggtgagcct	gtatcccacg	gggacgatgg	gctggggctt	3126
gttggttcaa	agaaactgag	aagtcagtct	tgggggaagg	ggtgggtagg	cctagaagaa	3186
aaaccgaaag	gagaccctga	gatctgcttg	aagttgggac	tctaaaagcg	tgtgtatgtg	3246
tttggtggtg	gagtggatag	aagtgcctag	aaatgtcagg	gtcgaggagg	gaacaggggt	3306
gggccagtcc	aggccctttc	agaattaagg	ctgggctaga	ccggttcctt	cagtatctcc	3366
ccagcctcct	tcctaccgt	gtacaatgca	ggggattatg	ctgaggggcg	gggccgcccc	3426
agtggccgtt	tggcccgtg	cacgtataaa	ggtgctacag	aaatgtgtag	tcgttcaaag	3486
tcccagggcg	ggcagacagg	caggcaggca	ggcgggcggg	cgggcgggcg	ggcgggcctg	3546
gggaggctct	gcatctgcag	ttgccgggga	agaataaagg	ggggaagagg	caagcactga	3606
tttacagtgg	ttcagtgcag	cttcaaaaaa	aaagggggaa	aaagcaatgg	gacttctggc	3666
tcagagtact	gctttgcatc	tcgtcttgca	taaatttgca	tactgagttc	aaagtcgtaa	3726
aatagactag	ggaaatgagt	gggatctact	gtcactgatt	gttaacacca	ctaccttcca	3786
cctcccctcc	ccccatctgg	acagaactcc	cggcactagt	tctgctaagc	agcgctgcct	3846
gagtcaccag	tcaccactgt	tcatgaaaga	aggagatggg	gaagcgcctg	tctaggagac	3906
caagcagggg	gtggcagtgt	gcaccctggc	ctcggaggca	gggcctctct	gaaggggtgtg	3966
gcccagacaga	ttctggacca	gtgaggcaag	tggtgttctt	tctgacgcct	ggggctgccg	4026
ttgcccattg	tgggctgcag	tttgaggaag	ctgcagaatc	cgggtcttct	ttgtttgaaa	4086
tttccctact	cagcttccac	atatgtccaa	aacatttact	gtgataagtc	cctttgtgtg	4146

- 21 -

gtggtaatga	cccacttctt	tagacaagat	at ttgaggaa	ggttcgagaa	cagaaaaatt	4206
ctacttaaca	atgttaagga	atctgagaaa	agtggggcac	aggagtctct	aggttagaat	4266
aggtaggtaga	gcctggaagg	agagaagaac	ccatctgttg	tattcccagag	ggtctgcccc	4326
cacaatggct	cttggccaag	tggactgttg	agcttcctgt	caggcagagg	gaagagggag	4386
gcttagatag	ttctcactgt	tggcctactt	cagcactgag	cttcaagact	ttcaggcaaa	4446
tcattccttc	aggctcagag	gctggaaatt	agttctataa	aaatgccatg	gggccggtcg	4506
cattggctca	tgcttgtaat	cccagcactt	tgggaggccg	aggcagggtg	atcacaaggt	4566
caggagtttg	gagaccagcc	tggccagtat	agtgaaacct	cttctctgct	acaaatacaa	4626
aaattagctg	ggcatgggtg	agcatgcctg	tagtcccagc	tgcttgggag	gctgaggcag	4686
gagaattgct	tgaacccggg	aggtaggagct	tgcagtgagc	tgggattgca	tcactgctct	4746
ccagcccggg	caacagagca	aaaaaactcc	gtctcaaaaa	ataaataaaa	taaaaatgcc	4806
acgggttggc	caggcgcggt	ggctcacgcc	tgtaattcca	gcacttaggg	aggctgagg	4866
gggcagatca	cctaaggcca	ggagttcaag	accagcctgg	ctaacatagt	gaaaccccat	4926
ctctactaaa	aatgcaaaaa	ttagttgggc	atgggtggcac	ctgcctgtag	tcccagctgc	4986
tcgggactct	gaggtaggag	aatcgcttga	acctgggaac	tggaggttgc	agtgagccga	5046
gatggcgcca	ctccactcca	gcctgggcaa	caagagtga	actccgtctc	aaaaataaaa	5106
taaaataaaa	ataaaaataa	aaagccatgg	gttacctgaa	agtaggagca	tatccattca	5166
ctcattatct	gacctttgag	ggctgacttg	gagagaactg	ttctctggcc	cagtcatcag	5226
gtggcttcca	gttgggttctc	agctgggctg	ttcacaccca	tagaaaatgg	gaggtacgac	5286
tgttcaattt	cttttttttt	ttgagacgga	gactcgctct	cttgcccagg	ctggagtgca	5346
gtggtgcat	ctcggtcac	tgcaagctcc	gtctcctggg	ttcacgccat	tctcctgcct	5406
cagcctccc	agtagttggg	actacaggca	cctgccacca	cgcccggcta	actttttgta	5466
tttttttagta	gagacgggg	ttcacagtgt	tagccaggat	ggtctcgatc	tcctgacctc	5526
atgatccgcc	cacctcgcc	tcccaaagt	ctggggatta	caggcatgag	ccacgggtgcc	5586
cgaccaggct	aatttttttt	tttttttttt	tttttgagac	ggagtcttgc	tccgtcgccc	5646
aggctggagt	gcagtcgccc	agcctggagt	gcagtggcgt	gatcttggct	cactgcaagc	5706
tctgcttcct	gggttcacgc	cattgtcctg	cctcagcctc	ccgagtagtt	gggaattcac	5766
gcatgcacca	ccatacacgg	ttaatatattt	tgtattttta	gtagaggcag	ggttttacca	5826
tgttggccag	gttgggtctca	aactcgtgaa	cctcaagtga	tctgcccgcc	ttggcctccc	5886
aaagtgctgg	gattacaggc	atgagctacc	gcacctggcc	acaactgttc	gatattgata	5946
tgaggaagtt	ctgaaaattc	tgcaccaaag	ggaatggaga	acaggtaa	acctatgatg	6006

- 22 -

cctggcagga	taggacaggg	ggctgctgaa	aggaatgaag	gaggagcaga	acaaaaaggg	6066
gctgtgaggc	cacatccaca	tctgcttttg	ttgttggtcc	ttcccgcgta	ctagaagact	6126
gtacccaaaa	gaaaagaaaa	aagcttggtt	tttttttggt	tgtttttttg	tttttacaaa	6186
gagtctgtgt	atctgggaat	tgttattttt	atttgtatta	ttatttttat	tcatttatatt	6246
atttcaagac	cgagttttgc	tctgtcacc	aggctggagt	gcagtggcgc	aatctcggtt	6306
cacggcaacc	tctgcctccc	gagttcaagc	aattctcctg	cctcagcccc	ccaagtagct	6366
ggaattatag	acacctgcca	ccatgcctgg	ctaatttttg	tatttttagt	agagacaggt	6426
tttcaccatg	ttggccaggc	tggtctcgaa	ctctggacct	caggtgatcc	tcccgccttg	6486
gcctcccaaa	gtgctgtgag	tacaggcgtg	aaccactgca	cctggcctat	tttaattttt	6546
taaggaactc	tgtatccagg	gagtgcattt	cgtctcagcc	tttctctgta	tgttcccgac	6606
tccctcttcc	tcctcctgcc	ttctccatgc	tgteectctt	gtccctgcag	accattcttt	6666
cctttcccac	acttctccct	tccatgcctt	tcctactccg	gctcagctag	aacacctgct	6726
tgctaggtag	gtgggtgggg	ctcaaaaatg	gtgtggaaaa	tatagtggtc	tcagcctcag	6786
agatccatcc	aatcagaggg	atattatgta	ggtgactgga	gactataatg	cttgaaaaaa	6846
ctagaatcct	taaatgagga	gggtggttga	ctgggctttg	agggaaagga	cttctatagg	6906
ccagggaatg	ggattctctc	ctataagagt	tagcactagc	gctgcttctg	ctgaaaccaa	6966
ggcatagaag	cgggagatga	tgtttgaatt	catgagggaa	gctggtaacg	tggttggcag	7026
atattctaga	tccctctgct	tatcctgtct	ctggccagca	attcactgag	atggcaggca	7086
gccgccagga	gattggtggg	gctcgcactc	gtcctcacc	tattatgaac	tggttctgt	7146
gcctgctttg	tctcctgtct	tccacctccc	tgcgagggag	gccgttctac	ctccattttg	7206
caggtgtgga	cactgggtga	ctcagcgata	cgaaaatgct	gagaggctaa	cctgggtacc	7266
ttcccactgc	accctctac	ctccgtggcc	gggtaaggag	ttgggctggt	ggtagagaaa	7326
agcttttgac	cgtatcaacc	agcagttttg	ttcctggaca	gctgccatgt	gcaggtgatg	7386
tatggggcac	agtaaatttc	tgccagattt	caattatgta	gattcccttg	actccatgga	7446
cagttacaag	tcctggagtt	tggggacctc	acagttgtaa	gtagccagca	tacagaggtc	7506
agtcaagagg	gacaagaagg	tcacaggggtg	ggcctcttga	gagaagagct	gctgggtagc	7566
agatgatggg	gcaggcagcc	tggcctttgc	tgtcatattg	accgggggtg	gttccattgc	7626
aaccctgtct	cttaataact	cagtagttag	taaggctctt	cctccaatgc	cagttccctg	7686
ttggtaaaat	agcaatgatt	ttacctcttt	cccagggtgc	gtgttggggg	ggctgtgata	7746
atggcaagta	cctggcacat	agtatgttct	tgtacatggt	agctgcctta	atcttggcaa	7806
gtgacatctc	ctccttgaca	attagctggc	ctttaaaatg	aaggccaaac	tggaaagggtc	7866

- 23 -

aggcaggcca	ctcctctctg	gaggttcaaa	tcccagggtg	tctaccatgt	actaatcacc	7926
caaggtaaga	gatttcttgc	taagcctcag	ttttcttatt	tgcaaatacag	aaattgatag	7986
tatttcatg	ggtcaactac	acagatgtgt	gtaaagatat	aaactatata	taaaacctaa	8046
gctcttgcat	tttggtgatg	agtcacttga	tttccatctc	ttcacctatg	aatggaaata	8106
tccatacctc	tccaacctac	tttcacaggg	ctgtagtgag	gtgggtttatg	gaaagggcaa	8166
tgaaactgta	ctttaaatgc	aatacaaatg	caattgataa	atgcaatata	aatgcaactg	8226
ataagtgc	tataaatgca	gtgcataaac	ggttggagac	catcgggctt	gcattttttt	8286
tttttttttt	tttagacaga	gtcttgctct	gtcgcccagg	ctggagtgca	gtggcgcgat	8346
ctcagctcac	tgcaagctcc	acctcctggg	ttcacgccat	tctcctgcct	cagcctccc	8406
agtagctggg	actacaggca	cccgccacca	cgcccggcta	attttttgta	tttttttagta	8466
gagacgggg	ttcacagtgt	tagccaggat	ggtctcgatc	tcctgacctc	atgatccgcc	8526
tacctcagcc	tcccaaagtg	ctgatgatta	caggcgtgag	ccatgggtgcc	cgaccaggct	8586
tacatttttt	tttttttttc	gtgagacgga	gtctcgctgt	gtcgcccagg	ctggagttca	8646
gtggcacgat	ctcggctcac	tgcaagctcc	atctcccggg	ttcatgccat	tctcctgcct	8706
cagcctcctg	agtagctgga	ctataggcac	ttgccaccac	gccagctaa	ttttttgcat	8766
tttttagtaga	gatgggggtt	cacagtgtta	gccaggatgg	tctcgatctc	ttgacctcat	8826
gatccgcccc	cctcggcctc	ccaaagtgct	gggattacag	gtgtgagcca	ccccgtctgg	8886
ctaggcttgc	attttttttt	tttttttttg	agacggagtc	ttgctctgtc	accaggctg	8946
gagtgcagtg	gcgogatctc	ggctcactgc	aagcccgcct	cctgggggtt	atgccatttt	9006
cctgcctcag	cctccccagc	agctgggact	acaggcgcac	gccgccacgc	tcggctaatt	9066
ttttgtat	tttagtagaga	cgggggtttca	ccgtgttagc	caggatggtc	tcgatctcct	9126
gaccttgtga	tccgcccgc	tcggcctccc	aaagtgctgg	gattgcaggc	gtgagctacc	9186
acgcctggcc	taggcttgca	tttttaaatg	attagttggg	cagtgtctct	tggatccgat	9246
gtgctgatgt	ctaggactga	ctgttttttg	tatgggagct	tctgtctgca	gtgtctagaa	9306
tgggggttgg	agggactatg	gggaacagag	gtggccccag	agcttggaca	tctgggttct	9366
catgctggcc	ttgctgctca	taagcaggta	acacagggtt	gagttcctcc	tttgccctggc	9426
cagccaacct	ggaagacct	gccttctggc	tgtcagtgtc	cagagctctg	gagaatccct	9486
gaggctacat	ctctgggtgg	gggaagggtc	tctctgtata	gatggccctc	caggggaagt	9546
ctgcaaggat	ccagacttca	gaataataga	gttaagtatt	aactaatcca	agtggtccta	9606
gcacatttag	cacatatggc	atgctcaagg	gtgtgcaatt	gtgcagtc	caaagccagt	9666
gagggcaccc	gatgccaccc	atttcctgca	ccaggaaaat	ggacctttca	cgtctctgaa	9726

- 24 -

cctgggatgc	cagaatgcac	tgaacttgat	cctggcaaat	aaattggaag	ctgtttccaa	9786
aaaagtgata	tgttcttgct	acattgtctc	cagagtttac	tcctgagttg	gggagcagca	9846
tctcattttg	caaggggctg	ttggattctc	catagaaacg	ctgggttggg	ggagtgtctt	9906
ctgagtactg	gggctctacc	taatgtctga	ttccatctgt	ggtaccaaaa	agccctctga	9966
atggcagcct	gtctgtgcac	aatgatgccg	aagactttct	ttccagcat	ctaatgaggc	10026
atggccccag	gaaagcagag	aaggcggatg	cttgagtctc	tgagggtcag	aatacgctta	10086
tggtgaaatc	agaactgggg	acaaggcaaa	gtactgccat	tgggaatccg	gggtggatgg	10146
tcatgttaga	ccaggggtgt	ccaatctttt	ggcttccttg	ggccacttgg	aagaagaatt	10206
gtcgtgggtc	acacataaaa	tacactaaca	ctaaccatag	ctgatgagct	aaaaaaaaaa	10266
aaattgcaaa	aaaatctcat	aatgttttaa	gaaagtttac	gaaattgtgt	tgggccacat	10326
tcaaagctct	cctgggctgc	aggttggaca	agcttggcgt	agactcttgg	agggctttca	10386
tggtggaggc	aggaaaggaa	aagccccctt	tacagatggg	atagtaggtc	cagagaaatg	10446
tgttggtttt	cagttccaga	gtaatttcac	ctgctccac	cctccacact	cacaaagagc	10506
aggctgggtg	ggatgtcgat	ttttttttct	tctggagaat	ggaccatctc	cctatgcctc	10566
tcctcctcct	tggtactaa	cccctgactt	ttccctcccc	cttctctggc	cactcctgtt	10626
cctgaacaaa	gccaggggaag	atctgggctg	gctcagcccc	tgggtgtcag	tcaagactca	10686
gagggagact	ttgctcaaaa	agccagccct	cccggaaaga	aatacagagc	aactgtagtc	10746
aggaattttt	ccctttcctt	ttttctgtgg	gcggggggag	ggacacttgg	cagtctttgg	10806
ggcccttctg	gctgccagcc	ccacatctct	tcctggctgt	tgggaggcag	ggcaggaggg	10866
aagattgggg	agaagggggc	aggggctggc	aaacacccag	ctgctgggca	gtggaattcg	10926
tccataggcc	ctgggtggagt	tggcctctag	tgaagcctgg	ggaggtttct	aattagtgga	10986
ggacggtgaa	ggtgtggcag	cctgtgttca	ttcaagggaac	tgagctctct	gagccaggaa	11046
gtcaaaggga	agagggggac	tgagtcccgg	aaggccttgg	taggtgggga	ggaagtgtga	11106
cctactcagt	gtgttccatc	ctattagaag	ttgagaagag	gggggtgttt	cagctggagt	11166
gattcaggtt	ggactctggg	ggtgcagaga	gtcagtcatt	ggcactaagc	aaagggagcc	11226
acctcaaaga	ataagtgata	taacagtgat	aaacaaaatg	gctttgttct	agacattgcc	11286
tgcttctacc	agaaagctca	ttttgttttt	tttgtttgtt	ttatatctct	cccctctccc	11346
acttcctgcc	ccagctctgt	tttagcctcg	agattctctc	cacttaacaa	atgaagaaat	11406
caaggctaca	ttcaggttca	actgccaaag	caagcacaga	ccagattgta	gtctgggtct	11466
ctagattctt	ccagctctcc	atttcaccag	actcacacat	gcagaaaagc	agtgaattat	11526
atactgcttc	ttggaagagc	tccttttccc	tgggaagtag	taattaaggc	atatcttctc	11586

- 25 -

aaaaatgttc	ctttatgcaa	ctctctagga	agtttctttt	aggtctattg	tctttacttc	11646
tggatggagg	tgccaagttg	cctctgggtg	gtgaacttgt	tttctaaaat	gcccgaggcc	11706
aagaacaaaa	tgaaaatggc	aattcccttg	tgccctgtgc	cctgccccgt	gacttcctga	11766
taatgctgag	cagttgctat	agcaacagtg	gtgggaggtg	tgtcaggcag	gataaagggg	11826
gtgggggtag	gacaggttta	ctggatcaaa	gaggcttttg	cggttgagta	gggctgctga	11886
catgactgat	agcctacttc	tagggaaaga	tgggggtccc	atagagatgg	ggaggggaag	11946
aatgaagcc	ctggggctaa	aaggtcttgc	ttgagagtct	ttccttaggt	gatgccaagt	12006
gaccagggtgc	catctgggtg	gttaacaacc	tgggcttggt	tttggccact	ggctacatga	12066
ctttgtgcag	ttggcctcca	gctctatagt	tcatttatca	agtgtgaggg	ggataagaaa	12126
ttaccaacct	catgtgcctg	cttgataaca	aaggtaaaca	tgcaagccac	gtagcacaga	12186
aactcagata	atcaaatgag	gtagtgcagg	ttctgattct	tgcaatacag	gagaacagga	12246
gaatggcaga	caccgctttt	ggacgttttc	tgattggtea	tttcaagaat	cctgttagca	12306
gggtgtgactt	attcccttga	gtgtggctta	cctggacagc	tgccactttg	ctttggaaag	12366
cctccagtga	gtccagagag	gagtcaatgg	ttgagagttt	tccagggggc	agtcatgacc	12426
ctcgatgcca	ttgtgcctgt	ggttctacca	acagcttaat	gaccaaccat	gtcccattta	12486
tgccccagaa	gttaggactc	ccacttgagg	ctctgctggg	tgtctctggg	ttggttcact	12546
gcctcttaaa	ccttgagagc	atagccaagg	aaggtaattg	atgtgatata	ctatcagaca	12606
aaattagtga	acttttagttg	aagacagaga	aaagcatggg	gacctattta	aagacactta	12666
tttatattca	agacacttct	tgtgctgggt	tgagagtggc	cggcaatcct	ggacatcttt	12726
gcaggtttta	aatttctttt	tgcacccctg	gggttcaatt	tgtcaggcct	ggaatatctg	12786
agagcagagt	tatggggatg	gagaagagga	cagtgggaagg	ggaggggtgg	gggaaagggt	12846
ggtaggattc	agcctttgtt	acactatggg	agtggggctg	tgggattgca	tcatcgaaac	12906
tgaattgttt	gaatgtgggt	cttaggacag	gcatgtgtcc	ttcccatccc	atctcatggg	12966
tccacttgct	cactgccttt	cttgacttgc	tcactgcctt	tcttgactct	taaccactct	13026
tgctgatacg	ggctctagat	gaggcaaagg	ccagaagaat	gagaaaaaga	gttaatgtaa	13086
cagggtactc	ttgggcaagt	gctagatgct	ctttgatagt	gtattagtct	gttttcacgc	13146
tgctgataaa	gacataccca	agactgggta	atttacaaaa	gaaagagggt	taatagactc	13206
acagttccat	gtggctgggg	aggcctcaca	atcatgggtg	gaggtgaagg	cacgtctcac	13266
atggtggcag	acaagagagc	ttgtgcaggg	aaactcccct	ttataaaacc	atcagatctc	13326
atgagacttg	ttcactatca	cgagaacagc	atgggaaaga	cctaccccca	tgattcaatt	13386
acctcccact	gggtccctcc	tacaacatgt	aggaattgtg	ggagctacaa	ttcatgggtga	13446

- 26 -

gatttgggtg	gggacacagc	caaaccatat	cagatgggtca	cctttttctaa	tgtgtaaaca	13506
ttaattgagc	acatgctgtg	tgtcagctag	gagaccatat	gaaaaagcag	ttgtcatacc	13566
acctgtaaca	catgcttttt	catgctggag	ggggacaaga	aaagcatttt	agaaaggtga	13626
ataaagaaga	aaccaaggca	gcagagatgt	atatacatcc	tatgcaaata	aacaggtagc	13686
tcttggtcac	taaaagagct	taaccaagtc	tcttgggaag	agcttgatgt	taaataatga	13746
agtccttca	tggtaaataa	agttatgaaa	gtaagagaat	caagttggac	aggcttctac	13806
ttcttcagtc	ttggaagaaa	aaaaaaagca	gtgtcttcct	tcctaaaatg	ttaattgagg	13866
acacagaatg	tcttctcacc	tgtccttcat	tcactaagta	tggaaagctg	tacggaagct	13926
aagaccaagt	ttatgtcttc	aggaagcttc	tgtccttagg	aggactaaga	aatgcacag	13986
acctacagct	ttttggcaat	ggactcaggc	tagagtgtg	tgaagaaagt	tgggtcaa	14046
taccagaaga	ctttaccag	cagtgtcttt	cttttttaaa	attttgtaaa	atttttattt	14106
atttatttat	tttatttatt	ttgagacaga	gtccccctgt	tatgcccagg	ctggagtgc	14166
gtggcgtgat	ctcggcttac	cgcaagctcc	gcctcccggg	tttgcaccat	tctcctgcct	14226
cagcctcctg	agtagctggg	actacaggcg	cccaccacca	cgcttggtta	attttttgta	14286
tttttttagta	gagacgggg	ttcaccatgt	tagccaggat	ggtctcgatt	tgctgacctc	14346
gtgatctgcc	cgccttggcc	tcccaaagt	ctgggattac	aggggtgagc	cactgcgccc	14406
ggcccttatt	tatttgtttt	tttgagacaa	gctctggctc	tgtcacctag	gctggaatgc	14466
aattgcacga	tctcagctca	ctgcaacctc	cacctcccgg	gctcaagcca	tcctcccagc	14526
tatggcagtg	cttttcatca	atgcaatcaa	gtgacctttg	gacacttgga	ggccatattg	14586
ttcatccttc	ttccttctca	aaaggtttct	cctaagttat	accagaggga	gtctgtgcaa	14646
tttcttccca	tctgcaattt	atcagatttg	ttttgttctt	tcttatgaaa	ttcttggatg	14706
gcttctaact	tggatctctc	ctgttttagt	ttaaggctgt	tttctctctg	gtccttgggg	14766
agacagaacg	gctccttccc	ttcccttggg	ttttcctggc	ccttggacac	ctccaccact	14826
cagcaccctc	tggcagcacc	aggatggatt	tgtcccttgg	ctttcagttc	tcaggcctga	14886
gtcatcttga	ttacttcac	ttcccccttct	taattctgtg	accctcttct	actacctttg	14946
ggtcttcttt	ctgactcatt	tccgagttct	tcatggcata	gagaacaggc	caggagccca	15006
gggctagggt	ctgactgggtg	gacaagtgtg	cctggtaaca	tcattgccct	ggccagtgtg	15066
agtgggtgggtg	aagtccgcca	tgtcactatc	gtttgggtcct	cttccagtag	ggcttcaggg	15126
acacttgagg	ttactgatag	ctctgcttga	tgaggaagat	gaggggtgggt	tgggtgtgaga	15186
ggtgactttt	ttgtttttgt	ttttattttt	gagacagagc	ctctcactct	gtcaccagc	15246
ctggagtcag	tggcgcgatc	ttgggtcact	gcaaccttcg	cctcccgggt	tcaagcgatt	15306

- 27 -

cttctgcctc	agcctcctga	gtagctggga	ttacaggcat	gcaccaccat	gcctggctaa	15366
tttttgtatt	tttactagag	acgggggttc	accatattgg	ccaggctggg	ctcgaactcc	15426
tgacctcgtg	atccacctgt	cttggcctcc	caaagtgctg	ggattacagg	catgagccac	15486
cgcgcccagc	cctgagaggt	gactttttta	aaagaatatt	taaaaaatgc	agaaaagtac	15546
aaagaacaat	gtaataacac	aagtctacca	cccagaaccg	tgggcgggtg	atattttgtc	15606
cactagcacg	cagtacttac	ttagtgttta	tctttcttag	ctaagcagac	ttgcatatat	15666
gtgatagatg	actaatatgc	ctatgtttgc	atatattgca	tctatatgca	tatatatgat	15726
agatgaccaa	tattgtccaa	attgttttgt	gaaaataaat	tgaaaatgca	cataaagggt	15786
ttattttccc	tcttaacatt	aggtgtctga	gatctaattg	tgacatccat	aggtcagaaa	15846
tattcttttt	aactgctata	tattgttcct	ttgtattacc	ctactgggtt	tatttatcca	15906
ttcacctggt	gaaacacttt	tggatctggg	tgctaatagt	gtttttacaa	ctacaagtaa	15966
tgctgtaatg	accatttttg	tacatgtgtc	cctgtgcaca	tgtagagaga	agaattgctg	16026
ggtcaggcac	attcttccca	ttcccatagt	ccctgggttc	agtttgaatc	aaagcacagt	16086
ctactcactt	gggcctcagc	aactagggtc	ccttgccagg	ttcctggatg	gctttagttg	16146
aaccaagga	gaaatgtttg	ggacccgtgg	aaaggagtga	aagctgatag	tgctgtttgc	16206
catcttgata	tcagtttctc	tggctcttca	aggacagctt	ggctcacggg	gccccagagg	16266
accttggctg	aatttgagca	tgtttttggt	ttctcaaact	gtctgcctca	gccctgaccc	16326
tccccaggtc	tttgcaattc	caggaatgac	cctgatgttt	ataaaggagc	cctttggaga	16386
gtgcacaggc	agggaggctc	tggagaggct	gtgctgtttc	tgttcactcg	gggggagccc	16446
gagatgagat	ggcagcatga	gggcatactc	tagaaacctc	catgacgcca	ggccaggaga	16506
ggccctctg	catgtatcca	agcaagcata	tgtgtctgtt	tacaaaccca	tgagtggatg	16566
agcttacaag	gctgtgtctc	tgcaaagctt	atacccgaga	ccacctctct	tgaaaagaag	16626
aggcattggt	ctcgcctcac	caggaagtcc	ttcaggaaac	agacagaaca	gagagaatca	16686
caaaggcatg	ggggcctggc	aatgccagag	cttgagagga	gaggtatcat	tcaacataca	16746
gattccttag	tccttgctc	tgactttctg	gggctggggg	ccagggtctt	tatttttagc	16806
aagctcatca	gatgatttct	gatgaagagc	tatgtttggg	aacttctgac	tgggtctaata	16866
tccctgaatt	tacagatgag	tacactgaca	cctgcgcaag	ttgaacaccc	tgcttagcgt	16926
cactcagaat	cccaggggtg	gtgggtgaagt	caggatcgaa	cctcagtcct	tcctcccaca	16986
ctgggactgc	ttggccaagg	cttcgaaccc	aaaagctggg	gctccaaagg	aaatccaaaa	17046
agagtttgca	gagtggtttg	acgtggaggg	gattcatcgg	ccagggtggg	gctctctgcc	17106
ctcccttggt	ggtgcatcat	ctcctgtagg	tgttttgtca	ccgtggaagg	gcagtagcct	17166

- 28 -

gtgcctgttg	ggcttgccac	tcaacagagg	tcactgagct	cttccgcact	cagctgggtg	17226
attggcctgt	ggtagctgca	ggtggaggga	gatgattctc	agaagagaaa	ggccagttct	17286
tgcagtcgtg	tgatcttata	acctagggga	cttgtggatc	tgaaacatct	gagaaggctg	17346
tttgcagtta	tttactacca	caaaggaaac	tgggagagga	gggaccacct	gttggaaaac	17406
catgctcaga	agtgactggg	ctgttgctga	aggatggaag	gatatcaggt	gttgcattgg	17466
tggctccctt	gcaggaagga	agggtcagg	gggtgagaag	gaataggaag	ggagtattaca	17526
atgaatgaag	ctgggtgtcc	tcaggagata	gatggaccag	ttgggggtgat	gtcattaact	17586
aagctggaat	gtgtcctgag	tgccagccta	taccgcgaca	gacatctgtg	cacactattc	17646
agaattccag	tgtgagaggc	cactgcccc	tctgaccttt	tgaaaaaaaa	agtcagggga	17706
agacacataa	acagtgtcac	tcacccacaa	agtctcagga	cctcagggtta	aacttttagct	17766
aaagaagtaa	caaatagaac	ccccatttta	ccttcctctt	tcctaacttc	tgcctccctc	17826
cctcccacca	gcatggcctg	aattactaat	cccagcctgt	tggtgggctg	tgctattgaa	17886
catagtggcc	acatgctaga	tggagtgttg	tgaccaccat	cagccatcca	aagaggactg	17946
gcttcaaatt	ctctccaaca	aaagctgttt	cttgtatgtg	ctggcgtggt	tatgagccct	18006
gtctcaactg	agcatgcgtg	gtgattatga	gctactgcct	gacctccttc	cctttgtccc	18066
tcccgctaaa	ttctgtaaac	aaagttaaag	taatttaggc	aaattcattc	ttgcttaagg	18126
acttctatta	aatggcagag	agctttgata	tttagcaatg	tataaaattt	gtacaatcaa	18186
aaccaagctt	tagaagaaat	atgcaataga	tcattattgca	gaactctcat	gactcatttt	18246
ttccatgcat	ttggccattc	tacaaatcag	acattaccta	gccttcccct	cacatggcag	18306
atataacttt	tatttaccca	ttcaagaaat	tgattgttca	cacattcctt	tttctttgaa	18366
acaaggctct	gctttgtcgc	ccaggctgga	gtgtagtggt	gtaatcttgg	ctcactgcaa	18426
gaagtgaact	ccgcctgctg	ggttcaagcg	attcttgtgc	ctcagcctct	caagtagctg	18486
ggacctcagg	catgtgccac	catgcccagc	taatttttgt	atttttggta	gagacagggt	18546
tttgctatgt	tggccaggct	ggtcttgaac	tcctggcctc	aaatgacctg	cccacctcag	18606
cctcccaaag	tgctgggatt	acaggcatga	gccaccatgc	ctggccacca	cacatttttt	18666
tctaaagctt	ggttttggcc	acagtgagag	tttcttgggc	tgtcaggggt	aaaaatacca	18726
tgtcttgttt	aacagcagcc	atcatttttag	cagaattttt	tttttttttt	tttttgagac	18786
tgagctttgc	tcttgttgcc	caggctggag	cgcaatggcg	cgatcttggc	tcactgcaac	18846
ctccgcttcc	cgggttcaag	tgattctcct	gcctcagcct	cctgagtagc	tgagattaca	18906
ggcatgcgcc	accatgcctg	gctaattttt	atgtattttt	agtagagacg	gtgttactcc	18966
atgttgggtca	gactgggtctc	gaactcctga	cctcagggtga	tctgcccgcc	tcggcctccc	19026

- 29 -

aaaatgctgg	gattacaggc	gtgagccacc	gcgccccgcc	ccttagcaga	attgttaata	19086
tcattttttca	gacttgagaa	attagagctc	agagaggtta	aaggttctga	ctcgtgagtg	19146
aatctcaaag	ggcaaagctg	gcattcaggg	ttcttgactg	ctgggttcat	tgctcattcc	19206
agtttggttaa	tgaatcgtcc	actgctatgg	atattaatgg	agtcgtctag	agagcagggg	19266
gaggaggatg	gtggttagta	tttattctgt	tgcccaacat	atgtcaaggc	taagtgcctt	19326
ttgggagaaa	ttaacaaaga	cggagccatg	tgagttttct	gcccagctag	ggtggaacca	19386
ggaagctgcg	gccagccttc	attcccctgg	cccattactc	agaattaact	cttgaaaggc	19446
ttggaatgct	attacctgtt	gcaagaggtg	agttcattgg	aggaatgata	agttcagggg	19506
tggttagat	catctctgtg	gtactcctct	ttcccaaccc	tgctcttaga	ttttaagata	19566
atcatgagat	aggaggagaa	ggctggaaaa	tggttaagtct	ttcttcagaa	taaagaaaag	19626
aagaaagaag	agagacacat	gagctgttag	gaagatgtga	ttgttctagg	tagggaaaag	19686
gtcttatttc	ctcaccaggt	ctggataagg	accggggcca	cacactctga	ggaggaagga	19746
ccagggaggt	gcttttagat	tcttggggct	tagtttccta	agaaagcttt	ttaaggtctc	19806
tcttagatca	ggaatcacca	atttccaaat	agagcactga	ggtcagaaca	ttgtgctagg	19866
agaaaatgag	atthttgtggg	agtcctgttc	tatcaaacac	aagagtttga	ttgtgtgagt	19926
aagatcattg	atgcaacaaa	caggattaca	ttggaaaaat	gagctgtgcc	aaaattaggt	19986
ctgaaaacct	gggtatgaat	gcatagaagt	gaacaataaa	ggctctttct	cttgggctgt	20046
cagcttccag	ctgctctcct	tggctaaatt	agtaagggtg	ggccaaaaga	caatgtaggg	20106
accccatgat	aattgtgaag	aatggcagag	aggttgtttg	ctcctgtgaa	ttcaaattgg	20166
agtccatctg	ggctcaggtt	tgccatctgt	aaaatgggac	aacgatccat	ttcttttcat	20226
cccatctgcc	ttcctcatct	ccagctctcc	tctaggggtc	ttaggtagaa	aggtgtggca	20286
gtctacccaa	ggcaggggtg	ggagttctcc	cctgcccttc	aagccccga	ctaatggctc	20346
tgggggtgtc	tgctcatctc	ctgacttctc	tttggaagaa	cttgaccttc	tatatgtgaa	20406
atcccttgct	tctgtgttga	cccatttccc	taggtgctat	gtagggaaga	agagctaatt	20466
cttactaatc	tacagggaaa	gtctcttcag	gggccttgga	ggtttctatc	aaccatttct	20526
aagcccatga	gccttttcca	ttgatgtcat	ttgggggttg	taagaacat	ttcttatagt	20586
gacctttgca	ccatagccca	tggagttgca	tacataagt	gttgcttaat	aaatattatt	20646
ctgaacttaa	ctctcttagg	cagaaaaaaa	atacaacttg	gtatthttgt	tctgagggca	20706
aattccatct	ctagctggag	ttcgacatct	ggaatgtttt	cctgtggatt	gttcttgcag	20766
cctataagag	gatatatggg	aagaacttgg	taacatgtga	taatgccctt	gtctgactga	20826
ggggaggagg	ggcgctatgt	gggtacttcc	tggtaggcag	gaaatggaag	tggctaagag	20886

- 30 -

ctaacccecat	gtgtgcctac	tgtgtgccag	gccctctctg	tagatgcagg	cagcatctct	20946
ttggatcctc	ccagttgccc	ataggaggtg	catactgttt	gtcctttttac	agatgaacaa	21006
tggaagcaca	gagagcttaa	gaaagttgtc	tggtgctgca	cagctcagta	gtaaaagctt	21066
gaattccaac	tcaggcagtt	agatttttggg	gtttgaatct	ctttggctgg	gaacagtaac	21126
aaggaatgca	ggctttttctc	agcctacaat	gctttctgat	ctctgcctgg	tcaactcttt	21186
catttccctt	ccctgggctt	ctacttcaag	atgtggagtt	cagatctggc	tacgggattt	21246
gagtccagaa	gcctcttcct	gagaaagaaa	cagatgctga	tgcattggtct	tgttaaacca	21306
tctttaaaag	accttgcatc	agcaaaccag	gcctgtggat	ttcttcatct	tattggcatg	21366
ttcgttggca	ctgctgagat	caataaagaa	atggaatgaa	acataccctt	gaatccaggt	21426
ggcctctgtg	agggtggaca	aacttgacct	tcacctcaga	gagtttggga	catctcaggg	21486
caggaaatca	actttgtcct	cctggaggaa	cactgccaa	ttgaccccag	cctcctggga	21546
cctggagcca	gggttgcata	tgacatctgt	gggccctgga	cactgcctct	gtgggcccctt	21606
ctataaaaac	atatattttt	aaaattttaa	aattattatt	tttcgttttt	gagacagggg	21666
ctcactctgt	caccaggtct	ggagtgcagt	ggcatgatca	tggtctactg	cagcctctaa	21726
cccctgggct	caagtaattt	cccacctcag	cctcccaggt	agctgggacc	acaggtgtgc	21786
atcaccatgc	ctggctattt	ttttattttt	agcagagaca	aggtgatata	gttaggcttt	21846
gtgtcgccac	ccaaatgtca	tcttgaattg	taatcccat	aatccccagg	tgttgatgga	21906
gagatcaggt	ggaggtaact	gaatcatggg	ggtgggtccc	ccatgttggt	ctcgtgatag	21966
cgagtgagtt	ctcctgagat	ctgatgggtt	tataaggagc	tcttccccgc	ttcacctggc	22026
acttcttcct	gccaccttgt	gaagaagggt	ccttgcttcc	cggacctttg	gcttctacca	22086
cgattgtgag	tttcctgagg	cctccccgcc	atgctgaact	gtgagtcaat	taaagctctt	22146
tcctttataa	attacccagt	ctcaggcagt	tttttatagc	agtgtgaaaa	tggactaata	22206
cagtcaggcg	tggtggctca	tgccataaat	cccagcactt	tgggagaccg	aggtgggcct	22266
cctgtggcaa	ggagttcaag	accagcctgg	ccaacatggt	gaaatcctgt	ctgtactaaa	22326
aatacaaata	ttaaccaggc	gtggtgacac	atgcctgtag	ttccagctac	tcgagaagct	22386
gaggcaggag	aattgctcaa	accaggagg	cggagggttg	agtgagccaa	gatcgtgcca	22446
ctgtactcca	gcctgggtga	cagagcgata	ctctgtctca	aaaaaacaaa	aaaaaaaaaat	22506
caaaaaacca	tattttataa	cttgtttaat	aaatgcaaat	gtattaatat	tatatagtaa	22566
gactttttta	cctaaaagtt	attttttttt	tctgatgtta	atagaaataa	aaatatgtcc	22626
taggcactgt	gcctccagag	cccaatggac	aagttggccc	tgccctggcg	cttcaggccc	22686
gaatgcacat	ttatctgtgt	gtgcccctct	cctttttcaa	atgtgtttcc	agagattgca	22746

- 31 -

gagggccttc	ttctgaggat	taatgttgga	ttgacctctg	ggacgcagcc	aagggatggt	22806
tctttggcca	ctgtctgtgg	tacaagaggg	cttcgtgaag	atggggagct	gcaggctgaa	22866
agtcctctag	gaagcccagg	gaagatcctg	tgccctgccc	ccaagaaact	gtggactatt	22926
gtggctgcag	atggaagggg	agcggagggg	tcttagccgt	ggatgcggaa	tatagtggag	22986
ctagatttgc	tttatatggg	gcctggagcc	atttcagcaa	tgggctatca	gcttacaggt	23046
cttgggctaa	aagcctgact	ccagaagagt	gattctcacc	ctgggcagtt	ttgcctctcc	23106
ggggccattt	ggcaacgaat	gaagacacat	tttttgttgt	tccaacctgg	ggaggggtgct	23166
actggtttct	ggtgggtaga	ggccaggaat	gctgcccac	atctggagta	cactggatgt	23226
tccaggcatt	gagttcgggt	gagcggctcc	ttcccagaag	ctcatggata	gaggtacaca	23286
tgagctgcct	ttcccgctct	caaccagaca	acgtgttggt	atagccataa	tagagactgt	23346
ctgccttcac	ccaagccact	tccaagagg	gcagagccag	gtgagcctct	gagcatcaca	23406
gacttcatgt	ccacagattg	gagagttcca	gaaatgggtg	caggggtgcag	taccagcaga	23466
ctcccagagc	tctgtcttct	tctaacaccc	tgtgagcaga	atgcgggtgtc	gaggtctcca	23526
gaaacccttc	ctgcctctgt	gggcacaaag	ctgggtaggg	tgcggtaagc	agggatgatgc	23586
caagggatgat	ctcaggtacc	aaggggtcgg	gtgagggcga	actggaggtt	cccacttcat	23646
tttgtcttct	tcattttgat	cctgagtagc	tgggagggag	tcaaagggct	cttctctatg	23706
ttccaatgca	ggtgggtggga	gtggggagac	tttctagcta	ctttgctctc	ctcctgtggt	23766
tttggtgtct	gcagagggca	ggggcagggg	gcttgagatg	gtcccactca	ggtctggggg	23826
cagagaggcc	tgtctgggaa	gggcagcggg	gcggaggcag	ggaggctggg	gatgagtgcg	23886
tgcgctgcac	ggcatctgtg	tgctgcgtta	gaggtcagtg	ttgagatgga	aagttgactg	23946
aacctgatct	cttctgtttg	ttgtccaggt	ttgtctgata	tcagactttc	cccggccttt	24006
ctctttcacc	cttttgtttc	tgacctctgg	ggtccatctg	aagaggggag	gggggctctt	24066
ctaggagggg	gtggacctgg	tggcgctgaa	gaaatctgcc	gcacgggtggg	gtgactcagc	24126
gtccctgttc	ttgggaacct	ggcaggcagc	tgagcagggg	cactcactca	gcctgcctct	24186
ctgggtccct	gggtgtctgc	tgctctccat	gtggaggagc	acagagggca	gggcaggatg	24246
gatgaggaaa	gctgtgggct	ggtgcctggg	tctgattaga	catggtggag	caggactcca	24306
acctagcctg	gaggcgcacg	ctaaatcggc	cttttcctgc	cggtgggtgct	gggctgccgg	24366
gctgctggaa	caatttaatc	tgtgctggct	tcatttatgg	tacaaccag	gctccccaga	24426
gcccctgtgg	gaataaccat	tctaaataaa	atatttttcc	aggtttcagg	tttatacatt	24486
taacatttaa	attttttttt	tcctggagtt	caacaatttt	ggataaaagt	cttttaaaat	24546
gatgaatagg	tagaatttta	attccctccc	ctttttgatg	tccaagatc	atcgatcatca	24606

- 32 -

acgtaattct	tgttttgggc	tgtaagacat	ggaaatcccc	agggttcatt	gaggggcatt	24666
cattgtttgc	cacacacaaa	atgtacccgg	agtttgtatg	ttttgttttc	ttagtgtcct	24726
ctttcctctc	tgccatttga	tccttcttgc	tactgtccac	atgcattcct	tctttgtccc	24786
ttcaattttg	ttgactttct	tctgcggtgg	ccttggaac	cagacatcca	gaagccaggc	24846
ttcctggagc	agcactgttg	tgaactgtct	cgaggctcct	cacgagatga	ttctcccctc	24906
atcttgagtg	gaaagtgcc	acatttgaac	atgctcagcc	gtggtaacag	ctattaggat	24966
gttcagagct	tattttcctt	ggtttccttg	ttctcggttg	ggttccatgc	tgctgaagta	25026
gtagataact	ggggcattcg	tcattgtcat	catcatcagt	gtaataacca	gcagcaacat	25086
ggatatagaa	tttctgtgct	ccaggctcgc	ttttatttat	ttatttat	atttat	25146
ttatttat	ttgaggcagt	gtcttgctct	gtcgcccagg	ctggagtaca	atggtgcgat	25206
ctcagctcac	tgcaacctcc	acctcctggg	ttcaagcaat	tgtcctgtgt	cacccttctg	25266
agtagctggg	attacagggtg	tgtgccacca	caccggcta	atttttttgt	atttttagca	25326
gagatgggg	ttcaccatgt	tggccaggct	ggctttgaat	tcctgacctc	aagtgatcca	25386
catgcctcag	cctcccaaag	tgctgggatt	ataggcatga	gccaccacac	ccggcctaag	25446
tcacttttta	actgcttttc	tttcgtcata	ctctcaacag	ctctatgaag	aaagaactgt	25506
tattatccag	caacttacag	ttaagaaagt	agaggcacag	agagggtctg	tgtggggaga	25566
acacgtgcag	ctgatataga	aaatggcaga	gctggacttt	aaccggcta	cacccttta	25626
accagagctg	ctaccctggg	gacggcgtgc	tggcttctaa	aggaagctgc	tgcattggcaa	25686
gtgaagagt	actagtctc	aggcttgag	ggaggggaaa	ggacagggac	agaccccat	25746
caaagtgtcc	gtatgctggg	gcctatgctg	ggcacgcctt	gacagcaggt	gctgctggag	25806
tgccagcacc	cgggtgctgc	cctggcatgg	aagagggtgc	ccctccctta	gttggtgagt	25866
gagtgaatac	tcttgtatag	agttactcta	agaattat	atttgtgttc	taagggtgt	25926
aagtttttga	aggtgcatta	atggggaaga	aaaaatgtgc	aagattgggt	gaaaaacacc	25986
cctctgcacc	caccagatc	ttgttgggac	ctgcaggaga	gcagctctgg	gtgggagggt	26046
ctggggctgt	ttgccgtg	tccaaggggt	tggggctggg	gctcggcgg	gtgaggccct	26106
gcccttctgg	agcttgctgg	ggcttgtctc	ctggttacag	ctctgccgtg	cgattcagat	26166
cctctctgcc	tcctttggct	ggacgtgcct	cagccagagt	gagactgtgg	caggctggag	26226
ctgttttcaa	ggcctccttc	acctaggctg	tgcatacaaa	aggttgaaga	aagtataagg	26286
agacttaatg	gacgcttgtt	ggacagacat	cacgaagctt	catgccctgc	agtaagctga	26346
agctatgaag	tcgaaggagt	catagttcct	gttcttgggg	ctctcagtat	aatgggggtg	26406
gggtgggctg	cacagacaga	cttgggaacg	agggaatgtc	agtgtggcag	agctgtagct	26466

- 34 -

ttgtccaggc	tgggtctcaaa	ctcctggact	caagcaatcc	tcccacctcg	gcctccccgaa	27957
gcgctgggat	tacaggcgcc	agccactgcg	cctggcctga	aactcagtct	tgttgccata	28017
tgcttctctc	actgagctac	ccctccccctc	tttcctgccc	agctctgcca	tcttgggtcc	28077
tggccctggg	tggttttatg	gaaagaagt	gagcttgggc	atgggggtggg	ggcatggaga	28137
ctcgagacgg	ccacttagat	ggatatttgg	aggcagagcc	cattctgaca	catgaaagcc	28197
acggtcacag	aacaaggaga	ggagtgaaca	ttcccatcca	tacaattttc	tttaggggct	28257
gggggttggg	acccagcatg	gcagggagat	ccccgggcag	gctgtcagaa	gtccctgtgc	28317
cccctgctgg	ctgtgtgacc	atgggaaagt	cccctccctg	tgcagcagct	tctcgctgag	28377
cttgcgctcg	gcgcgtgctt	cgtgggcctg	ctgtcctcac	tccggggcgtc	ccctgctgtc	28437
tgtgcgctgca	g act ccc tgc caa cag gaa ctg gac cag gtc ctg gag cgg					28487
	Thr Pro Cys Gln Gln Glu Leu Asp Gln Val Leu Glu Arg					
	225		230		235	
atc tcc acc atg cgc ctt ccg gat gag cgg ggc cct ctg gag cac ctc						28535
Ile Ser Thr Met Arg Leu Pro Asp Glu Arg Gly Pro Leu Glu His Leu						
	240		245		250	
tac tcc ctg cac atc ccc aac tgt gac aag cat ggc ctg tac aac ctc						28583
Tyr Ser Leu His Ile Pro Asn Cys Asp Lys His Gly Leu Tyr Asn Leu						
	255		260		265	
aaa cag gtgagcatgg tgccagcctg ggccagagca gctcctcctc agccctgggc						28639
Lys Gln						
270						
cccagatgtg ccttgtttct gccccacccg cctatgatcc tctgaggtct gagctgagtg						28699
agagactcag actcctgtca ccatggggat tgggatgcca gctgccaggc cggggagggt						28759
gcagctcttc tccctgcctc cgacgggtca gttgctggct ccacttgccc agctgcccac						28819
cggcctcact tggagtccag gttgggagga ggctcttgga gaagggaagt ggcttaaacc						28879
aagcatcttt gctttgatcg gacattgaca tcacctggag ctcttcctgg gcccgcccc						28939
aaattgccat gtggtaggtc aggtttacag ctctaacaag cctcagggtga ggctcataat						28999
ctgtgagcca cactcccacc atcattgatt tcaaccactg gccgagttgt cttcatgggt						29059
tcgaagtatc tcagggaatg aagagtcagg catggagaaa tgccgttgga aagattctga						29119
aaagccatct agaacaccat tgtaaataca ttggagaaaa tgaagtttaa gtaaaccact						29179
caggagccct tctccctcct ctatccacct tctgttcata tgcatttttaa cagtcactgt						29239
gcatatgcaa tattttccct gtaacatact tttttttttt ttttttttga gacggagtct						29299
cgctcttgtc tcccaggctg gagtgcagtg gcgtgatctc agctcactgc aacctccgcc						29359
tgccgtgttc aagcaattct cctgcctcag cctcctgagt agctgggatt acaggtgcct						29419
gccaccatgc ctggctaatt tttgtatttt tagtagagat ggggtttacc atgttggcca						29479

- 35 -

```

ggctggtctt gaactcctaa cctcagatga tccccccacc tcgacctccc aaagtgctgg 29539
gattacaggc atgagccacc acacctggct gggcattggg atttttataa agctaccctg 29599
gtggttctaa tggcagtctg ggctgacctt aagtgtggat gtgtgaagcc aggagatggg 29659
aacagggctc tgctgatggc ttaggggtct cacctctcca ccccttcac ccaggcacct 29719
tgacctcta aatattagtt tagtattaaa agctatttgt gtacatgccc tctaaatgta 29779
agctggctaa ctccatttgg agctcagaag aaaaagtctc gatcacttct catgggaaac 29839
aggctgtctg ctgcaccttt gggtaactaa agtgatttgc aggagtagtt agcttatgca 29899
taaaacagag accttggagt gggtaatgcct ggcttccatc tcagctgtgc ttcttaccag 29959
ctgtgtgacc ttgggcaagt tgcttagccc atctgtgctt cagttttatc atctgtaaac 30019
cagggataat aatggcatct aattcacaaa gtggttgtga ggattaaatg actatatttg 30079
ttagaaactg aactagcagt gtgtcttgca cacacagtgc ctggcacaca ctgattttga 30139
ttgcaacatg aggttttaca cctgacgcta tgactaagct tcagtatgct ggggtcccccc 30199
cactggatgg gacctctttg aggctgggga ctgagtcttt tcatcttcgg gtgtcctgaa 30259
tggccttgct aatcagaggg gtccctgggc gggcagcatc agcatcacgt tggagctcgc 30319
tagcgatgca gaatttccag ctctccaggg acctactgca tgggaatctg cattttatca 30379
aggtttcctg gcgcatattt gaagttgcga agctccaccc ggcagcgcac gggtagctgc 30439
aaggggtggc tgctcagtgg acgccgggtt gagggacaga aggaaagttg ctggctgcgg 30499
gctcctccat gctcttctcc tctctcccca g tgc aag atg tct ctg aac ggg 30551
                                Cys Lys Met Ser Leu Asn Gly
                                275

cag cgt ggg gag tgc tgg tgt gtg aac ccc aac acc ggg aag ctg atc 30599
Gln Arg Gly Glu Cys Trp Cys Val Asn Pro Asn Thr Gly Lys Leu Ile
    280                                285                                290

cag gga gcc ccc acc atc cgg ggg gac ccc gag tgt cat ctc ttc tac 30647
Gln Gly Ala Pro Thr Ile Arg Gly Asp Pro Glu Cys His Leu Phe Tyr
    295                                300                                305                                310

aat gag cag cag gag gct cgc ggg gtg cac acc cag cgg atg cag 30692
Asn Glu Gln Gln Glu Ala Arg Gly Val His Thr Gln Arg Met Gln
                                315                                320                                325

tagaccgcag ccagccggtg cctggcgccc ctgcccccg cccctctcca aacaccggca 30752
gaaaacggag agtgcttggg tgggtgggtgc tggaggattt tccagttctg acacacgtat 30812
ttatatattg aaagagacca gcaccgagct cggcacctcc ccggcctctc tcttcccagc 30872
tgcagatgcc acacctgctc cttcttgctt tccccggggg aggaaggggg ttgtggctcg 30932
ggagctgggg tacaggtttg gggaggggga agagaaattt ttatttttga acccctgtgt 30992
cccttttgca taagattaaa ggaaggaaaa gt 31024

```

- 36 -

<210> 10
 <211> 325
 <212> PRT
 <213> Homo sapiens

<400> 10

Met Leu Pro Arg Val Gly Cys Pro Ala Leu Pro Leu Pro Pro Pro Pro
 1 5 10 15

Leu Leu Pro Leu Leu Leu Leu Leu Leu Gly Ala Ser Gly Gly Gly Gly
 20 25 30

Gly Ala Arg Ala Glu Val Leu Phe Arg Cys Pro Pro Cys Thr Pro Glu
 35 40 45

Arg Leu Ala Ala Cys Gly Pro Pro Pro Val Ala Pro Pro Ala Ala Val
 50 55 60

Ala Ala Val Ala Gly Gly Ala Arg Met Pro Cys Ala Glu Leu Val Arg
 65 70 75 80

Glu Pro Gly Cys Gly Cys Cys Ser Val Cys Ala Arg Leu Glu Gly Glu
 85 90 95

Ala Cys Gly Val Tyr Thr Pro Arg Cys Gly Gln Gly Leu Arg Cys Tyr
 100 105 110

Pro His Pro Gly Ser Glu Leu Pro Leu Gln Ala Leu Val Met Gly Glu
 115 120 125

Gly Thr Cys Glu Lys Arg Arg Asp Ala Glu Tyr Gly Ala Ser Pro Glu
 130 135 140

Gln Val Ala Asp Asn Gly Asp Asp His Ser Glu Gly Gly Leu Val Glu
 145 150 155 160

Asn His Val Asp Ser Thr Met Asn Met Leu Gly Gly Gly Gly Ser Ala
 165 170 175

Gly Arg Lys Pro Leu Lys Ser Gly Met Lys Glu Leu Ala Val Phe Arg
 180 185 190

Glu Lys Val Thr Glu Gln His Arg Gln Met Gly Lys Gly Gly Lys His
 195 200 205

His Leu Gly Leu Glu Glu Pro Lys Lys Leu Arg Pro Pro Pro Ala Arg

- 37 -

210	215	220
Thr Pro Cys Gln Gln Glu Leu Asp Gln Val Leu Glu Arg Ile Ser Thr 225 230 235 240		
Met Arg Leu Pro Asp Glu Arg Gly Pro Leu Glu His Leu Tyr Ser Leu 245 250 255		
His Ile Pro Asn Cys Asp Lys His Gly Leu Tyr Asn Leu Lys Gln Cys 260 265 270		
Lys Met Ser Leu Asn Gly Gln Arg Gly Glu Cys Trp Cys Val Asn Pro 275 280 285		
Asn Thr Gly Lys Leu Ile Gln Gly Ala Pro Thr Ile Arg Gly Asp Pro 290 295 300		
Glu Cys His Leu Phe Tyr Asn Glu Gln Gln Glu Ala Arg Gly Val His 305 310 315 320		
Thr Gln Arg Met Gln 325		

<210> 11
 <211> 10884
 <212> DNA
 <213> Homo sapiens

<220>
 <221> promoter
 <222> (102) .. (1905)
 <223>

<220>
 <221> mRNA
 <222> (1906) .. (2440)
 <223> Exon 1

<220>
 <221> mRNA
 <222> (5726) .. (5952)
 <223> Exon 2

<220>
 <221> mRNA
 <222> (6497) .. (6616)
 <223> Exon 3

<220>
 <221> mRNA

- 38 -

<222> (8212) .. (10775)
 <223> Exon 3

<220>
 <221> CDS
 <222> (2038) .. (2440)
 <223>

<220>
 <221> CDS
 <222> (5726) .. (5952)
 <223>

<220>
 <221> CDS
 <222> (6497) .. (6616)
 <223>

<220>
 <221> CDS
 <222> (8212) .. (8334)
 <223>

<400> 11
 ctgcagacct gggacctcaa gaattgcatt tgatgccgaa cccagctcta atttcagagt 60
 caaggtctct gcgagtattt aaggaacgga tgtaaacctg ggggattcgt tttgtttcct 120
 tcaattttcc aatgaaatca gagatcctgt tcttgggtgt caacgcagat actagaagga 180
 ggtgatacaa gagaaaggaa acagcaagcg acgattatgg cacggtttcc tgtaaacaag 240
 gttgagtgta gccacagcct gagcactgtg ggagaagagc tcataagaaa atgacgggtgc 300
 tgggccttcg tcaccccggg gccctccatt gttcttgtct ttggtctctt tttatttgta 360
 gaggtccaat tatttattta tttagtacaa gagggaacga aattgatctt tccattctaa 420
 aaggagagta tatatgtata aaaggaagct gtatagatat gggggaagag gtggacaggg 480
 ggaaaagggg agaggacgag agagagaaag ggaggagagag ggacaaggag agacactggg 540
 cgagagatcg attaggagag acagaaatga tgaatgaaga ttaacttcac ccaaggcttc 600
 gtcgctggag gggaatggag gagctcctga tttgctatta ctactccaaa ctgcaaaggg 660
 ctccttcaag tcacctatcc acctcctaag gcaagcgtcc aatttcaaca gcgttcagga 720
 aagtctcttc ccgcggaggt ctaccgctt cccactccac cccacaaaac tctttggaaa 780
 agtgccttga aaaattttaat cctcaatcca atcctggacc accagcgtcc tctgttggtc 840
 accgaaggag ggggtgcgca gacaaaactg aagaaactcg agtgccagag aaggccgaca 900
 ggagttacag cgacctcagc gcgcaattgc gccccgaact ttactgaaaa gtgttttagat 960
 tgcagagata agctagaatc ccaacgcacg gagaatacag taatacgaag tcgccttcaa 1020

- 39 -

```

aaaatgacaa tgaaaattgc ctattaaagg actatttggt taattacgtt tcagcagtgc      1080
ccagttttatt gtcttttatta ttctttttgtc gtgggtgtaa actccatttg aaaacataat      1140
cagggagaaat acccaagaca agaagaacag ttgtcattta aaatatttga aaagccctgc      1200
cttaaggagc attcgcttgc cgggccactc ttaattgggg acttgcggtg tagcaacacg      1260
tgagagtctt cttgcgttga gaagtaagcc tggaaaggcg aaggccccgg ggcattctca      1320
gatgcgtatt tgtgggcccc tggggatata aacagcccag cgggtgtaaa ttaaaccctg      1380
cagtgccttg gctccctgag acccaaattgt aagtcagaaa tgtcccaaga cttcgccctgc      1440
caacggaatt aaattttaga aagctccacg aggtacacac gaatgcggag cgctgtatgc      1500
cagtttcccc gacaccggct cgccgcaggg agacctcacc ccgagagcgg aaggggtaag      1560
ggcggcgggg tcaaggagat cgggggtgct gagttggcca ggagtgactg gggtgaccgg      1620
gggtgctgag gtggcctgga gtgccgggggt ggccggggcac accttggttc ttgtagacga      1680
caaggtgacg ggctccgggc gtgagcacga ggagcaggtg cccggggcgag tctcgagctg      1740
cacgcccccg agctcggccc cggctgctca gggcgaagca cgggccccgc agccgtgcct      1800
gcgccgaccc gccccctcc caacccccac tcttgggcgc gcgttcgggg gcgtgtcctg      1860
ggccaccccc gcttctatat acgggcccgc gcgcccgggc cgcccagatg cgagcactgc      1920
ggctgggcgc tgaggatcag ccgcttctg cctggattcc acagcttcgc gccgtgtact      1980
gtcgccccat ccctgcgcgc ccagcctgcc aagcagcgtg ccccggttgc aggcgtc      2037

atg cag cgg gcg cga ccc acg ctc tgg gcc gct gcg ctg act ctg ctg      2085
Met Gln Arg Ala Arg Pro Thr Leu Trp Ala Ala Ala Leu Thr Leu Leu
1          5          10          15

gtg ctg ctc cgc ggg ccg ccg gtg gcg cgg gct ggc gcg agc tcg ggg      2133
Val Leu Leu Arg Gly Pro Pro Val Ala Arg Ala Gly Ala Ser Ser Gly
          20          25          30

ggc ttg ggt ccc gtg gtg cgc tgc gag ccg tgc gac gcg cgt gca ctg      2181
Gly Leu Gly Pro Val Val Arg Cys Glu Pro Cys Asp Ala Arg Ala Leu
          35          40          45

gcc cag tgc gcg cct ccg ccc gcc gtg tgc gcg gag ctg gtg cgc gag      2229
Ala Gln Cys Ala Pro Pro Pro Ala Val Cys Ala Glu Leu Val Arg Glu
          50          55          60

ccg ggc tgc ggc tgc tgc ctg acg tgc gca ctg agc gag ggc cag ccg      2277
Pro Gly Cys Gly Cys Cys Leu Thr Cys Ala Leu Ser Glu Gly Gln Pro
65          70          75          80

tgc ggc atc tac acc gag cgc tgt ggc tcc ggc ctt cgc tgc cag ccg      2325
Cys Gly Ile Tyr Thr Glu Arg Cys Gly Ser Gly Leu Arg Cys Gln Pro
          85          90          95

tcg ccc gac gag gcg cga ccg ctg cag gcg ctg ctg gac ggc cgc ggg      2373
Ser Pro Asp Glu Ala Arg Pro Leu Gln Ala Leu Leu Asp Gly Arg Gly
          100          105          110

```

- 40 -

ctc tgc gtc aac gct agt gcc gtc agc cgc ctg cgc gcc tac ctg ctg	2421
Leu Cys Val Asn Ala Ser Ala Val Ser Arg Leu Arg Ala Tyr Leu Leu	
115 120 125	
cca gcg ccg cca gct cca g gtgagccgcc cgccaggtgc gctgcgtgca	2470
Pro Ala Pro Pro Ala Pro	
130	
gcaccgccac tggcgccgaa gggcctgggg gttgctgggt gccgctgcgg gagactccgc	2530
ttttcttctc actggagata atatgtgggg aaactgaagg cgctccggga aaggtgaagg	2590
cggtcgccga gggaccctcc ccagccggcc ctctacttgc tcgattctct aagtgcagag	2650
tacttgtaaa ttgcaaagcg ctttcagtga aaatgggtaa aggtttccgg agctgagggg	2710
agcgggtaccg atgttttagct gttggaaaga tcctggacac aggagattct cctcgccccg	2770
cacgggtgca cacggactgc aatcccaggg atgcttgggg atgggggggat ataggcggat	2830
ttggaccaag gaaggtgggt aggcacgttg taggaaatag tacctctctt ttaaaatact	2890
gactttgcac agccttttgg tttgcaaagc aatgtctagt cccggtatgt ccaaaaacaa	2950
gtaaagtgga ttcgggtttt gatatcttct gcggttgga aacctgaagc tgaaaaagaa	3010
gtaacttctt aaggttacct agcggccaca acagagtgtg ggtttgaact ccgcgtgcca	3070
ctttcagtag cataccattc ttacaactcg ggccaccct gcacctgcgc cgacctcaa	3130
caaacttcca ggtgcgtggg ggggtgcgggc aatgtggact aagtcaattt caatgacacg	3190
gcaagggaat tggaatcagt cctaggctgt ctcccttctt aatctgaaat gggggggggg	3250
aatgagatgt tgttaagggg agccccagaa gaggaaaaat gcaaacattt ggcagagtta	3310
ccctcttgct tagccactat cagtatcagg cagacagcga ctctggtaag ggcattcacat	3370
tgttccctta aaaaaaggag cgggggttgt ttaaattggat ttggcagctg ttctttcaag	3430
cattcttagc cagcctcacc tagttatatg agaaataaag ttcttgcctt gcacagctga	3490
aggctgggag aattctcccc atcctaattc ccccaactcc ccaacgatca cgttggacag	3550
atgtcactgg gcaggcccc atctagggct agcaggatga acagtccctt tataatttat	3610
gtagctgtag agttccacgc ccgggtgaag ttattttctg gctcggcaag gctggctctg	3670
ttcaccctg agaaatgctg gattcatgga aaggcaagat gcctgaaaca tacactggct	3730
ctggtcagct gttaaagctg ctggaggcat ttgtctctcg gggcaaagtt atgtcatttg	3790
ccaagtgtcg tacattattg tgcattttgg ggtattcaaa aagtgatctt agaaatactg	3850
atacacatcg tcattcttgg gcttttagcaa tcatcatgat taccacctta gtagcactgt	3910
agtatagggt gatgtgagtt ataagattat aaaaagatct aagtgacttc tagaatctat	3970
ttgacaaaaa aaggtaaatt ttcgacagtc aaaagtcaca attatctgtt gcttaaatag	4030
aactgttttg tcttcatgcc ctagtctgca gccagggcat taagaagaaa ccaaggaaat	4090

- 41 -

ttaagaaatt	actcaagggt	cttagaaaaag	aagtataaat	acgtttat	acatggttctt		4150
agagtatttta	cattcttagt	atctcttttta	tctcagtatt	tccttgaaaa	agaaagcaag		4210
ctaagattaa	aagaaattga	aaccaaatcc	tcgcaggtag	ggacctcctc	tgtgagggtc		4270
tgtgctggac	cctgggaatg	tgtgcttccc	aaggatatgaa	accccttggg	gaactttaca		4330
gcaggacctc	agtgagctgt	ttggcagggtg	aggaaactaa	gacccagaga	ggagaggggac		4390
tttcctaagg	ccctggtgag	tgacctgcc	gtagccactt	ccagggggaga	gcagagcatc		4450
tgcagccaaa	tcattgcagc	cccaggtagc	tttctagata	gactgtggac	cagatggggcc		4510
acctgagctc	cctgctaggg	ttacacatta	tagccctgtt	tgtgtagtag	agaaatttca		4570
tgactctcaa	ttgtggactt	aagccgatgc	ctccagacct	tggcatggtc	cacaggccct		4630
gggagcatgg	gctctgaatg	tagcctttga	tccccatagc	ggtcttacag	cccctccaag		4690
ttcattctga	agaaggaatg	gagtgagaat	cctggctgca	gatccagtct	tgaatttagt		4750
catatactta	aaattccaat	tcaactgtta	acattccagc	atccatttta	agcatcagac		4810
tttcttcatt	tagcactttt	tattataaaa	gggagatctg	ctggaggggg	atttctccta		4870
ccccacccc	accaggggaa	ggaaaagctc	tttggcactt	agaagtctga	gccgtgagtg		4930
attgtctgca	tccatgtgct	gctgtgttca	cccgggggtga	aaaggactca	4990		
cttaggcagg	caccagcaag	atgcacaggg	tctgtgtaga	ccttgagt	tagagatgta		5050
acgggggacct	agaaaacaag	ccaccaacat	gcttgcatga	ttctgagccc	ctgaggcaaa		5110
acgctttgca	ggtaataatt	cagttttccc	atctgagctg	gacaccaagc	tcttataagc		5170
gtgtttacct	ggtagcattg	aggacggtac	tgggtcaacct	tgggaattccc	ataagggctt		5230
gttacaactc	agactcgtgc	cgccactcca	gcgtttccgg	agtggagaat	gtgcatttct		5290
tccaagtccc	cgggctgccg	ctgctcccgc	gggtgggagg	accacacttg	gagttgactg		5350
caaaatttct	gagccggcgc	tgcagcagcc	tcccgtggct	caggtctgcc	ccctgccggt		5410
ggaagatgaa	gcatactgcc	ttcacctact	gaggggcact	gaagcgtttg	tctgccttct		5470
ttagttgcag	ctacttagga	agagcacctg	tcagattgac	tttcaaacag	ataacttctt		5530
gaggtagagc	aaccaccatg	tagtgagtag	tatgatggaa	taataacttca	tcgagggtatt		5590
taaaaaaaaa	acctcacttg	gattgccaac	taatattgtc	atttacatgt	gacctgggtg		5650
caacgttaag	attttttacaa	gactgtgata	gatattgatg	actctcatgt	gtttgtctct		5710
cttggggcg	taaag ga	aat gct agt	gag tcg gag	gaa gac cgc	agc gcc		5760
	Gly Asn Ala Ser	Glu Ser Glu Glu Asp Arg Ser Ala					
			140		145		
ggc agt gtg	gag agc ccg tcc gtc	tcc agc acg cac	cgg gtg tct gat				5808
Gly Ser Val	Glu Ser Pro Ser Val	Ser Ser Thr His Arg Val Ser Asp					
	150	155	160				

- 42 -

ccc aag ttc cac ccc ctc cat tca aag ata atc atc atc aag aaa ggg	5856
Pro Lys Phe His Pro Leu His Ser Lys Ile Ile Ile Ile Lys Lys Gly	
165 170 175	
cat gct aaa gac agc cag cgc tac aaa gtt gac tac gag tct cag agc	5904
His Ala Lys Asp Ser Gln Arg Tyr Lys Val Asp Tyr Glu Ser Gln Ser	
180 185 190	
aca gat acc cag aac ttc tcc tcc gag tcc aag cgg gag aca gaa tat	5952
Thr Asp Thr Gln Asn Phe Ser Ser Glu Ser Lys Arg Glu Thr Glu Tyr	
195 200 205 210	
gtgagagctt ttcctcttgt taaaggagga gggcaagacc tgccaagcct gggactcag	6012
agcctcttga gggcaattct tactcaacaa accccagcgc ctggctgatg ggtgggcaac	6072
ccctagcccc tctgtgccct acctctctcc tctccttaca taaagaatat tgaccctttt	6132
ggagaatctt atgaggatca agctgaaata acactcttaa aagcatatgg gatgtcataa	6192
agacctctgc agataatgaa aatattctca taaagatagt tttatttact tcatcctcta	6252
tgcttggtga cctgctattg gttccatgcc agcttctgtg ccttactctg ggaagagcaa	6312
aaaggagaca gggagtgatg gttagcttat tcgggggact ttcgtgctac atcagacata	6372
aggtatctga ggagcaaatt acaggtocca cttttggtag ttgtgcagca tcgtaagatt	6432
tttaaagcac acattctaga gtaaaaactg tgactctgtt gctctgggtcc ttcctgatcc	6492
ccag ggt ccc tgc cgt aga gaa atg gaa gac aca ctg aat cac ctg aag	6541
Gly Pro Cys Arg Arg Glu Met Glu Asp Thr Leu Asn His Leu Lys	
215 220 225	
ttc ctc aat gtg ctg agt ccc agg ggt gta cac att ccc aac tgt gac	6589
Phe Leu Asn Val Leu Ser Pro Arg Gly Val His Ile Pro Asn Cys Asp	
230 235 240	
aag aag gga ttt tat aag aaa aag cag gtgagtgagg tcctcagtgt	6636
Lys Lys Gly Phe Tyr Lys Lys Lys Gln	
245 250	
gttttcttcc tcttctgttg acacagagga gaaacccatg tcaccagcgc ccaggctctt	6696
gtggccatag ctctaactct gagcctgtgc agcaccagtg cccaggactt ggtgccagtc	6756
tcaggaggtc agaccaaggg ctgctttgac ttgttgctct gaggctgct atattggcca	6816
taatcctcaa ccctagtgcc tttccaccac ccgcttccca ctctgtcct ttcaatgggt	6876
caccacagg cggacaagat gctgcccagt ggcacccttt ataaactgca agtggacatg	6936
ttaacacatt tggtaatgct gcgtcaggga gtgacatttc aaacaactat tatagtcagt	6996
ttccaagaag tgtgacatga ggtcatacca caaaaaagct taccctgaaa tcccacaatc	7056
gtcccctttc ctactgatgc cttcccgata gtgagcagggt tgcaatatta agattttgaa	7116
aaggctgttg ctagatgttg gtgactcgtg tgtctctgtc tcccttgggc ttttcaagga	7176
aatgctagtg agtgggggga tgactgcagc atggccagct tggagagccc agccatcccc	7236

- 43 -

agcacatacc aggtgtctgt cttggcgtgg aggggatgga acttgaaatc agacactcgg	7296
tccatgctgg ggatggccag tctctccaaa ctggcatgtg gtcttcctcc gagtcactgg	7356
catttccta gaaagtccaa gtgagaagaa ggcatgagag tcatcaacat caaacaacag	7416
tcttttcaaa atctttatat tgcaacatag tcccattcct ggaaaaggaa tggagtgaga	7476
atcctggcta cacatcagcc ccaaagttag tcattgccta aaatcccaat taacctgaaa	7536
atgatcaaac aaatttaaga tatagtaata ttaagctgta ataaatatgc ttctataggc	7596
tttgtgttat gtgatggcac tatttcaatt ggctttctaa ttggacaatt gatactatgc	7656
tatctacaga attggccttt ggagacctaa gtgagccaca gtggcctcag ggtgaccata	7716
tactaggatt catagcagtg gccacagtca gaagcctaag ctttcctcca ttgccattgc	7776
tcgtttatac cacgtttctg tcaaagtcatt attcattcaa caaagtcata ctgagaaggc	7836
gtcatgtgag gctggatgtg ggctccaaag tcatagctgt gacattcgca ggcagcggga	7896
tgttctcagt tccacatttg gcagagaagt cagtcaagag gttctacaag ggctggtgtc	7956
caccttatac tcctagaaac acaaaactgc cccaccccc gctttcttgg agcaggaagt	8016
tacaccacaca cgcattgcaca ggcgcacact cagcgggcct aggcagcgtg gctcttgtgt	8076
tgccttagct gaaatttctg ttgtgctttc tcagcatagc agagtcacgc tggcaaacca	8136
tcatgcgcc tggccaccga cctgacacca gaccaggag cattcacttc tctgtcttct	8196
gtttctctcc cacag tgt cgc cct tcc aaa ggc agg aag cgg ggc ttc tgc	8247
Cys Arg Pro Ser Lys Gly Arg Lys Arg Gly Phe Cys	
255 260	
tgg tgt gtg gat aag tat ggg cag cct ctc cca ggc tac acc acc aag	8295
Trp Cys Val Asp Lys Tyr Gly Gln Pro Leu Pro Gly Tyr Thr Thr Lys	
265 270 275	
ggg aag gag gac gtg cac tgc tac agc atg cag agc aag tagacgcctg	8344
Gly Lys Glu Asp Val His Cys Tyr Ser Met Gln Ser Lys	
280 285 290	
ccgcaagggt gagtactcag gaggggcagc ctgggctcca gggcctcact gtccttggac	8404
cagcctcagg ggctgggcgt ggccactggc ctccccagg cttacagacc caggagctgc	8464
agctcagggc cagaaagagc aaagcaaata ggacagagcc ctcagaaggg tgcagggaga	8524
gggagacccc atcaacccaa ccaaacaagt gtggggaagg aggccggcca gtgcacctca	8584
gggacactct gctttatctc agatacctca cagcacctaa gctatcattc atccacacac	8644
aaagtgaaga ttttcaaagt taggccttac ccgtgagtct ggaggtcatt tatcttcaca	8704
gagaacgttt atcgcagact gctaagatac atgttctaata taagatgtga tgtgagaacy	8764
ctgaatgctc gttggagact cagttgaagt gcagcttttt ttctgtcaaa tatataatga	8824
atattctggt agtctgtggc taatataatt ttaataaagt taatttaaata ctgatagaaa	8884

- 44 -

aatgaaattt	taaacgataa	tttttagagaa	tgctattata	tccagtcttc	ttttttcttt	8944
taataaatga	gggaactatt	gggggaaagg	aataaataca	ttttctttca	ttttattaag	9004
acaaatttag	taagcagaag	aaatttgcac	gttttagttat	aagggtttct	tttttcctta	9064
caagttggaa	aaaataattc	taattttaagg	gtaactcttt	gacaatgaac	actgtgagca	9124
gcatctggta	ctcgttgctt	tgtttgaaaa	catgagttga	gaccccagcc	gcacttgcag	9184
cctagtgcc	ttagcctgca	ggctgtgctg	gatattctcag	ggcaagagtc	gagccctttt	9244
gattttgggg	ggattatttc	aatatatttg	ctttttcttt	ttgttttagt	taatgtggag	9304
ctcaaataatg	ccttattttg	cacaaaagac	tgccaaggac	atgaccagca	gctggctaca	9364
gcctcgattt	atatttctgt	ttgtggtgaa	ctgatttttt	ttaaaccaaa	gtttagaaag	9424
aggtttttga	aatgcctatg	gtttctttga	atggtaaact	tgagcatctt	ttcactttcc	9484
agtagtcagc	aaagagcagt	ttgaattttc	ttgtcgcttc	ctatcaaat	attcagagac	9544
tcgagcacag	caccagact	tcatgcgccc	gtggaatgct	caccacatgt	tggtcgaagc	9604
ggccgaccac	tgactttgtg	acttaggcgg	ctgtgttgcc	tatgtagaga	acacgcttca	9664
ccccactcc	ccgtacagtg	cgcacaggct	ttatcgagaa	taggaaaacc	tttaaaccct	9724
ggatcatccg	acatcccaac	gcatgctcct	ggagctcaca	gccttctgtg	gtgtcatttc	9784
tgaaacaagg	gcgtggatcc	ctcaaccaag	aagaatgttt	atgtcttcaa	gtgacctgta	9844
ctgcttgggg	actattggag	aaaataaggt	ggagtcctac	ttgttttaaaa	aatatgtatc	9904
taagaatggt	ctagggcact	ctgggaacct	ataaaggcag	gtatttcggg	ccctcctctt	9964
caggaatctt	cctgaagaca	tggcccagtc	gaaggcccag	gatggctttt	gctgcggccc	10024
cgtggggtag	gagggacaga	gagacgggag	agtcagcctc	cacattcaga	ggcatcacia	10084
gtaatggcac	aattcttcgg	atgactgcag	aaaatagtg	tttgtagttc	aacaactcaa	10144
gacgaagctt	atttctgagg	ataagctctt	taaaggcaaa	gctttatttt	catctctcat	10204
cttttgtcct	ccttagcaca	atgtaaaaaa	gaatagtaat	atcagaacag	gaaggaggaa	10264
tggcttgctg	gggagcccat	ccaggacact	gggagcacat	agagattcac	ccatgtttgt	10324
tgaacttaga	gtcattctca	tgcttttctt	tataattcac	acatatatgc	agagaagata	10384
tggtcttggt	aacattgtat	acaacatagc	cccaaataa	gtaagatcta	tactagataa	10444
tcctagatga	aatgttagag	atgctatatg	atacaactgt	ggccatgact	gaggaaagga	10504
gctcacgccc	agagactggg	ctgctctccc	ggaggccaaa	ccaagaagg	tctggcaaag	10564
tcaggctcag	ggagactctg	ccctgctgca	gacctcggtg	tggacacacg	ctgcatagag	10624
ctctccttga	aaacagaggg	gtctcaagac	attctgccta	cctattagct	tttctttatt	10684
tttttaactt	tttgggggga	aaagtatttt	tgagaagttt	gtcttgcaat	gtatttataa	10744

- 45 -

atagtaaata aagttttttac cattaaaaaa atatctttcc ctttggttatt gaccatctct 10804
 gggcttttgta tcactaatta ttttatttta ttatataata attattttat taaaatgttc 10864
 cctgctttcc ctttttagcaa 10884

<210> 12
 <211> 291
 <212> PRT
 <213> Homo sapiens

<400> 12

Met Gln Arg Ala Arg Pro Thr Leu Trp Ala Ala Ala Leu Thr Leu Leu
 1 5 10 15

Val Leu Leu Arg Gly Pro Pro Val Ala Arg Ala Gly Ala Ser Ser Gly
 20 25 30

Gly Leu Gly Pro Val Val Arg Cys Glu Pro Cys Asp Ala Arg Ala Leu
 35 40 45

Ala Gln Cys Ala Pro Pro Pro Ala Val Cys Ala Glu Leu Val Arg Glu
 50 55 60

Pro Gly Cys Gly Cys Cys Leu Thr Cys Ala Leu Ser Glu Gly Gln Pro
 65 70 75 80

Cys Gly Ile Tyr Thr Glu Arg Cys Gly Ser Gly Leu Arg Cys Gln Pro
 85 90 95

Ser Pro Asp Glu Ala Arg Pro Leu Gln Ala Leu Leu Asp Gly Arg Gly
 100 105 110

Leu Cys Val Asn Ala Ser Ala Val Ser Arg Leu Arg Ala Tyr Leu Leu
 115 120 125

Pro Ala Pro Pro Ala Pro Gly Asn Ala Ser Glu Ser Glu Glu Asp Arg
 130 135 140

Ser Ala Gly Ser Val Glu Ser Pro Ser Val Ser Ser Thr His Arg Val
 145 150 155 160

Ser Asp Pro Lys Phe His Pro Leu His Ser Lys Ile Ile Ile Ile Lys
 165 170 175

Lys Gly His Ala Lys Asp Ser Gln Arg Tyr Lys Val Asp Tyr Glu Ser
 180 185 190

- 46 -

Gln Ser Thr Asp Thr Gln Asn Phe Ser Ser Glu Ser Lys Arg Glu Thr
195 200 205

Glu Tyr Gly Pro Cys Arg Arg Glu Met Glu Asp Thr Leu Asn His Leu
210 215 220

Lys Phe Leu Asn Val Leu Ser Pro Arg Gly Val His Ile Pro Asn Cys
225 230 235 240

Asp Lys Lys Gly Phe Tyr Lys Lys Lys Glu Cys Arg Pro Ser Lys Gly
245 250 255

Arg Lys Arg Gly Phe Cys Trp Cys Val Asp Lys Tyr Gly Gln Pro Leu
260 265 270

Pro Gly Tyr Thr Thr Lys Gly Lys Glu Asp Val His Cys Tyr Ser Met
275 280 285

Gln Ser Lys
290

INTERNATIONAL SEARCH REPORT

International application No.
PCT/AU02/00558

A. CLASSIFICATION OF SUBJECT MATTER

Int. Cl. ⁷: C12Q 1/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

SEE ELECTRONIC DATA BASES

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SEE ELECTRONIC DATA BASES

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Medline, CA:IGFBP, insulin like growth factor binding protein, liver, hepato, HCC, cancer, tumour, tumor, neoplasm, carcinoma, malignant

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 99/46597 A1 (DIAGNOSTIC SYSTEMS LABORATORIES INC) 16 September 1999 Whole Document	1 - 18
P, X	WO 01/53837 A1 (DIAGNOSTIC SYSTEMS LABORATORIES INC) 26 July 2001 Whole Document	1 - 18
X	WO 00/69454 A1 (BOARD OF REGENTS, THE UNIVERISTY OF TEXAS SYSTEM) 23 November 2000 Whole Document	27 - 29

☒ Further documents are listed in the continuation of Box C

☒ See patent family annex

<p>* Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier application or patent but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p>		<p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&" document member of the same patent family</p>
---	--	---

Date of the actual completion of the international search
16 May 2002

Date of mailing of the international search report
12 JUN 2002

Name and mailing address of the ISA/AU
AUSTRALIAN PATENT OFFICE
PO BOX 200, WODEN ACT 2606, AUSTRALIA
E-mail address: pct@ipaustalia.gov.au
Facsimile No. (02) 6285 3929

Authorized officer
CRAIG ALLATT
Telephone No : (02) 6283 2414

INTERNATIONAL SEARCH REPORT

International application No.
PCT/AU02/00558

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	JG Reeve et al <i>Int J Cancer</i> (1992) 51(5) pp 818 - 21 "Insulin-like growth-factor-binding protein gene expression and protein production by human tumour cell lines" Whole Document	1 - 18
X	SA Weinzimer et al <i>J Clin Endocrinol Metab</i> (2001) 86(4) pp 1806 - 13. "Transferrin is an insulin-like growth factor-binding protein-3 binding protein" Whole Document	24
X	T Yamashita et al <i>Biochem Biophys Res Commun</i> (2001) 282(2) pp 647 - 54. "Serial analysis of gene expression in chronic hepatitis C and hepatocellular carcinoma" Whole Document	1 - 18
X	N Kondoh et al <i>Cancer Res</i> (1999) 59(19) pp 4990 - 6 "Identification and characterization of genes associated with human hepatocellular carcinogenesis" Whole Document	1 - 18
X	JG Scharf et al <i>Lab Invest</i> (2000) 80(9) pp 1399 - 411 "Analysis of the IGF axis in preneoplastic hepatic foci and hepatocellular neoplasms developing after low-number pancreatic islet transplantation into the livers of streptozotocin diabetic rats" Whole Document	1 - 18
X	Y Gong et al <i>Mol Cell Biochem</i> (2000) 207(1-2) pp 101 - 4 "The expression of insulin-like growth factor binding proteins in human hepatocellular carcinoma" Whole Document	1 - 18
X	M van Kleffens et al <i>Endocrinology</i> (1999) 140(12) pp 5944 - 52 "Generation of antisera to mouse insulin-like growth factor binding proteins (IGFBP)-1 to -6: comparison of IGFBP protein and messenger ribonucleic acid localization in the mouse embryo" Whole Document	24
X	Y Shirota et al <i>Hepatology</i> (2001) 33(4) pp 832 - 40 "Identification of differentially expressed genes in hepatocellular carcinoma with cDNA microarrays" <i>Hepatology</i> Whole Document	1 - 18
X	SG Gray et al <i>Br J Cancer</i> (2000) 82(9) pp 1561 - 7 "Altered expression of members of the IGF-axis in hepatoblastomas" Abstract, pp 1565 - 7	1 - 18
X	H von Horn et al <i>Cancer Lett</i> (2001) 162(2) pp 253 - 60 "Expression levels of insulin-like growth factor binding proteins and insulin receptor isoforms in hepatoblastomas" Whole Document	1 - 18
X	L Xu et al. <i>Cancer Res</i> (2001) Apr 1 61(7) pp 3176 - 81 "Expression profiling suggested a regulatory role of liver-enriched transcription factors in human hepatocellular carcinoma" Figure 5	1 - 18

INTERNATIONAL SEARCH REPORT

International application No.
PCT/AU02/00558

Box I Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos :
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☒ Claims Nos : **19 - 23, 25, 26, 30 - 37**
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
The claims are not limited to the technical features that define the invention.

3. ☐ Claims Nos :
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a)

Box II Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Claims 1 - 18 are directed toward a method for detecting the presence of liver cancer by detecting the aberrant expression of a gene encoding an insulin like growth factor binding protein (IGFBP).

Claims 19 - 37 are directed toward methods of treating liver cancer by modulating expression of an insulin like growth factor binding protein.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims
2. ☒ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/AU02/00558

This Annex lists the known "A" publication level patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent Document Cited in Search Report				Patent Family Member	
WO	99/45697	EP	1066716	SE	9800686
WO	2001/53873	FR	2803918		
WO	2000/69454	NONE			
END OF ANNEX					